



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IDS (m): 293 Lysate: sc-178771

BACKGROUND

IDS (Iduronate 2-sulfatase), also known as SIDS, is a 550 amino acid protein that localizes to the lysosome and belongs to the sulfatase family. Expressed in lung, liver, kidney and placenta, IDS uses calcium as a cofactor to catalyze the hydrolysis of select sulfate groups on dermatan sulfate, heparan sulfate and heparin and, via this catalytic activity, is essential for the lysosomal degradation of both dermatan and heparan sulfate. Defects in the gene encoding IDS are the cause of mucopolysaccharidosis type 2 (MPS2), more commonly known as Hunter syndrome, which is characterized by skeletal deformities, hepatosplenomegaly and progressive cardiopulmonary deterioration, as well as neurological damage and, in some cases, death. IDS exists as two alternatively spliced isoforms, designated long and short.

REFERENCES

- Wilson, P.J., Meaney, C.A., Hopwood, J.J. and Morris, C.P. 1993. Sequence of the human iduronate 2-sulfatase (IDS) gene. *Genomics* 17: 773-775.
- Malmgren, H., Carlberg, B.M., Pettersson, U. and Bondeson, M.L. 1995. Identification of an alternative transcript from the human iduronate-2-sulfatase (IDS) gene. *Genomics* 29: 291-293.
- Li, P., Bellows, A.B. and Thompson, J.N. 1999. Molecular basis of iduronate-2-sulphatase gene mutations in patients with mucopolysaccharidosis type II (Hunter syndrome). *J. Med. Genet.* 36: 21-27.
- Bonuccelli, G., Di Natale, P., Corsolini, F., Villani, G., Regis, S. and Filocamo, M. 2001. The effect of four mutations on the expression of iduronate-2-sulfatase in mucopolysaccharidosis type II. *Biochim. Biophys. Acta* 1537: 233-238.
- Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 30990. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Ricci, V., Filocamo, M., Regis, S., Corsolini, F., Stroppiano, M., Di Duca, M. and Gatti, R. 2003. Expression studies of two novel in CIS-mutations identified in an intermediate case of Hunter syndrome. *Am. J. Med. Genet.* A 120A: 84-87.
- Tomatsu, S., Orii, K.O., Bi, Y., Gutierrez, M.A., Nishioka, T., Yamaguchi, S., Kondo, N., Orii, T., Noguchi, A. and Sly, W.S. 2004. General implications for CpG hot spot mutations: methylation patterns of the human iduronate-2-sulfatase gene locus. *Hum. Mutat.* 23: 590-598.
- Parkinson-Lawrence, E., Turner, C., Hopwood, J. and Brooks, D. 2005. Analysis of normal and mutant iduronate-2-sulphatase conformation. *Biochem. J.* 386: 395-400.
- Sukegawa-Hayasaka, K., Kato, Z., Nakamura, H., Tomatsu, S., Fukao, T., Kuwata, K., Orii, T. and Kondo, N. 2006. Effect of Hunter disease (mucopolysaccharidosis type II) mutations on molecular phenotypes of iduronate-2-sulfatase: enzymatic activity, protein processing and structural analysis. *J. Inherit. Metab. Dis.* 29: 755-761.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

CHROMOSOMAL LOCATION

Genetic locus: Ids (mouse) mapping to X A7.1.

PRODUCT

IDS (m): 293 Lysate represents a lysate of mouse IDS transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

IDS (m): 293 Lysate is suitable as a Western Blotting positive control for mouse reactive IDS antibodies. Recommended use: 10-20 µl per lane.


Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

IDS (B-5): sc-365047 is recommended as a positive control antibody for Western Blot analysis of enhanced mouse IDS expression in IDS transfected 293 cells (starting dilution 1:100, dilution range 1:100-1:1,000).


RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



IDS (B-5): sc-365047. Western blot analysis of IDS expression in non-transfected: sc-110760 (A) and mouse IDS transfected: sc-178771 (B) 293 whole cell lysates.



IDS (E-4): sc-365149. Western blot analysis of IDS expression in non-transfected: sc-110760 (A) and mouse IDS transfected: sc-178771 (B) 293 whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.