



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MFRP (m): 293 Lysate: sc-178932

BACKGROUND

MFRP (membrane frizzled-related protein) is a single pass type II membrane protein with two cubilin (CUB) domains, two LDL-receptor class A domains, a cysteine-rich domain (CRD) and seven N-glycosylation sites. The C-terminal CRD is related to the Wnt-binding domain of the frizzled family of transmembrane proteins, suggesting that MFRP may act as a Wnt receptor. MFRP is predominantly expressed in retinal pigment epithelial cells, co-localizing to the plasma membrane with CTRP5. MFRP interacts with CTRP5 via its CUB domain and may play a role in the development of the eye. Functional MFRP is necessary for photoreceptor maintenance and appears to regulate the axial length of the eye. Mutations in the gene encoding MFRP can affect various structures in the eye and may result in nanophthalmos 2 (NNO2), an eye disorder characterized by extreme hyperopia.

REFERENCES

1. Katoh, M. 2001. Molecular cloning and characterization of MFRP, a novel gene encoding a membrane-type frizzled-related protein. *Biochem. Biophys. Res. Commun.* 282: 116-123.
2. Kameya, S., Hawes, N.L., Chang, B., Heckenlively, J.R., Naggert, J.K. and Nishina, P.M. 2002. Mfrp, a gene encoding a frizzled-related protein, is mutated in the mouse retinal degeneration 6. *Hum. Mol. Genet.* 11: 1879-1886.
3. Jones, S.E. and Jomary, C. 2002. Secreted frizzled-related proteins: searching for relationships and patterns. *Bioessays* 24: 811-820.
4. Pauer, G.J., Xi, Q., Zhang, K., Traboulsi, E.I. and Hagstrom, S.A. 2005. Mutation screen of the membrane-type frizzled-related protein (MFRP) gene in patients with inherited retinal degenerations. *Ophthalmic Genet.* 26: 157-161.
5. Sundin, O.H., Leppert, G.S., Silva, E.D., Yang, J.M., Dharmaraj, S., Maumenee, I.H., Santos, L.C., Parsa, C.F., Traboulsi, E.I., Broman, K.W., Dibbernardo, C., Sunness, J.S., Toy, J. and Weinberg, E.M. 2005. Extreme hyperopia is the result of null mutations in MFRP, which encodes a frizzled-related protein. *Proc. Natl. Acad. Sci. USA* 102: 9553-9558.
6. Katoh, M. and Katoh, M. 2005. Identification and characterization of rat Ror1 and Ror2 genes in silico. *Int. J. Mol. Med.* 15: 533-538.
7. Ayala-Ramirez, R., Graue-Wiechers, F., Robredo, V., Amato-Almanza, M., Horta-Diez, I. and Zenteno, J.C. 2006. A new autosomal recessive syndrome consisting of posterior microphthalmos, retinitis pigmentosa, foveoschisis, and optic disc drusen is caused by a MFRP gene mutation. *Mol. Vis.* 12: 1483-1489.
8. Mandal, M.N., Vasireddy, V., Jablonski, M.M., Wang, X., Heckenlively, J.R., Hughes, B.A., Reddy, G.B. and Ayyagari, R. 2006. Spatial and temporal expression of MFRP and its interaction with CTRP5. *Invest. Ophthalmol. Vis. Sci.* 47: 5514-5521.
9. Shu, X., Tulloch, B., Lennon, A., Vlachantoni, D., Zhou, X., Hayward, C. and Wright, A.F. 2006. Disease mechanisms in late-onset retinal macular degeneration associated with mutation in C10TNF5. *Hum. Mol. Genet.* 15: 1680-1689.

CHROMOSOMAL LOCATION

Genetic locus: Mfrp (mouse) mapping to 9 A5.1.

PRODUCT

MFRP (m): 293 Lysate represents a lysate of mouse MFRP transfected 293 cells and is provided as 100 µg protein in 200 µl SDS-PAGE buffer.

APPLICATIONS

MFRP (m): 293 Lysate is suitable as a Western Blotting positive control for mouse reactive MFRP antibodies. Recommended use: 10-20 µl per lane.

Control 293 Lysate: sc-110760 is available as a Western Blotting negative control lysate derived from non-transfected 293 cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.