



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CCRF-HSB-2 Whole Cell Lysate: sc-2265

BACKGROUND

Santa Cruz Biotechnology offers a variety of whole cell lysates for use in combination with our antibodies as Western Blotting controls. CCRF-HSB-2 Whole Cell Lysate is derived from the CCRF-HSB-2 cell line using a procedure that ensures protein integrity and lot-to-lot reproducibility. All lysates are tested by Western Blotting to assure that each one contains the expected concentration and assortment of proteins. Numerous antibodies directed against a wide array of mammalian proteins are used to test each lysate. CCRF-HSB-2 was derived from the same buffy coat preparation as CCL-120 (CCRF-SB) by serially transplanting into newborn Syrian hamsters.

REFERENCES

1. M.D. Anderson Hospital and Tumor Institute. 1967. Research Report. University of Texas at Houston: 21.
2. 1967. Proc. Am. Assoc. Cancer Res. 8: 1.
3. Adams, R.A., Flowers, A. and Davis, B.J. 1968. Direct implantation and serial transplantation of human acute lymphoblastic leukemia in hamsters, SB-2. Cancer Res. 28: 1121-1125.

SOURCE

CCRF-HSB-2 Whole Cell Lysate is derived from the CCRF-HSB-2 cell line.

Organism: *Homo sapiens* (human)
Tissue: Peripheral blood
Disease: Acute lymphoblastic leukemia
Cell Type: T lymphoblast
Growth Properties: Suspension

PRODUCT

Each vial contains 500 µg protein in 200 µl of an SDS-PAGE Western Blotting buffer, which consists of 100 µl RIPA Lysis Buffer and 100 µl Electrophoresis Buffer, 2X.

APPLICATIONS

CCRF-HSB-2 Whole Cell Lysate is provided as a Western Blotting positive control. Recommended use is 50 µg (20 µl) per lane. Sample vial should be boiled once prior to use.

PREPARATION METHOD

Cells are cultured with appropriate media conditions and allowed to reach a confluency of 75%. Cells are lysed using the RIPA Lysis Buffer System (sc-24948). The BCA Protein Assay Kit (sc-202389) is used to determine the total protein concentration. The lysate is adjusted to contain 500 µg of total cellular protein in 100 µl before adding an equal volume of Electrophoresis Sample Buffer, 2X (sc-24945). Final concentration of product is 500 µg total protein in a final volume of 200 µl.


STORAGE

Store at -20° C; stable for one year from the date of shipment. Non-hazardous. No MSDS required. Minimize repeated freezing and thawing.


RESEARCH USE

For research use only, not for use in diagnostic procedures.

DATA



IL-2 (LNKB-2): sc-52017. Western blot analysis of IL-2 expression in Jurkat (A), CCRF-CEM (B), HuT 78 (C), H9 (D), SUP-T1 (E) and CCRF-HSB-2 (F) whole cell lysates.



Lck (D-8): sc-166628. Western blot analysis of Lck expression in CCRF-HSB-2 (A), Jurkat (B), Molt-4 (C) and HuT 78 (D) whole cell lysates.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.