



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

U-87 MG Cell Lysate: sc-2411

BACKGROUND

Santa Cruz Biotechnology offers a variety of whole cell lysates for use in combination with our antibodies as Western Blotting controls. U-87 MG Whole Cell Lysate is derived from the U-87 cell line using a procedure that ensures protein integrity and lot-to-lot reproducibility. All lysates are tested by Western Blotting to assure that each one contains the expected concentration and assortment of proteins. Numerous antibodies directed against a wide array of mammalian proteins are used to test each lysate.

U-87 MG cell line was established from a 44 year old male Caucasian. This is one of a number of cell lines derived from malignant gliomas.

REFERENCES

- Ponten, J. and Macintyre, E.H. 1968. Long term culture of normal and neoplastic human glia. *Acta Pathol. Microbiol. Scand.* 74: 465-486.
- Beckman, G., Beckman, L., Pontén, J. and Westermark, B. 1971. G-6-PD and PGM phenotypes of 16 continuous human tumor cell lines. Evidence against cross-contamination and contamination by HeLa cells. *Hum. Hered.* 21: 238-241.
- Fogh, J., Wright, W.C. and Loveless, J.D. 1977. Absence of HeLa cell contamination in 169 cell lines derived from human tumors. *J. Natl. Cancer Inst.* 58: 209-214.
- Fogh, J., Fogh, J.M. and Orfeo, T. 1977. One hundred and twenty-seven cultured human tumor cell lines producing tumors in nude mice. *J. Natl. Cancer Inst.* 59: 221-226.
- Olopade, O.I., Jenkins, R.B., Ransom, D.T., Malik, K., Pomykala, H., Nobori, T., Cowan, J.M., Rowley, J.D. and Diaz, M.O. 1992. Molecular analysis of deletions of the short arm of chromosome 9 in human gliomas. *Cancer Res.* 52: 2523-2529.
- Li, Y.M., Mitsuhashi, T., Wojciechowicz, D., Shimizu, N., Li, J., Stitt, A., He, C., Banerjee, D. and Vlassara, H. 1996. Molecular identity and cellular distribution of advanced glycation endproduct receptors: relationship of p60 to OST-48 and p90 to 80K-H membrane proteins. *Proc. Natl. Acad. Sci. USA* 93: 11047-11052.
- Clark, M.J., Homer, N., O'Connor, B.D., Chen, Z., Eskin, A., Lee, H., Merriman, B. and Nelson, S.F. 2010. U87MG decoded: The genomic sequence of a cytogenetically aberrant human cancer cell line. *PLoS Genetics* 6: e1000832.

SOURCE

U-87 MG Whole Cell Lysate is derived from the U-87 MG cell line.

Organism: *Homo sapiens* (human)
 Organ: Brain
 Tumor Stage: Classified as grade IV as of 2007
 Disease: Glioblastoma; astrocytoma
 Growth Properties: Adherent epithelial

PRODUCT

Each vial contains 500 µg protein in 200 µl of an SDS-PAGE Western Blotting buffer, which consists of 100 µl RIPA Lysis Buffer and 100 µl Electrophoresis Buffer, 2X.


APPLICATIONS

U-87 MG Whole Cell Lysate is provided as a Western Blotting positive control. Recommended use is 50 µg (20 µl) per lane. Sample vial should be boiled once prior to use.


PREPARATION METHOD

Cells are cultured with appropriate media conditions and allowed to reach a confluency of 75%. Cells are lysed using the RIPA Lysis Buffer System (sc-24948). The BCA Protein Assay Kit (sc-202389) is used to determine the total protein concentration. The lysate is adjusted to contain 500 µg of total cellular protein in 100 µl before adding an equal volume of Electrophoresis Sample Buffer, 2X (sc-24945). Final concentration of product is 500 µg total protein in a final volume of 200 µl.

DATA



ASRGL1 (LA-52): sc-130472. Western blot analysis of ASRGL1 expression in HeLa (A) and U-87 MG (B) whole cell lysates and human testis tissue extract (C).



DGUOK (H-3): sc-376267. Western blot analysis of DGUOK expression in U-87 MG whole cell lysate (A) and mouse kidney (B) and mouse thymus (C) tissue extracts.

STORAGE

Store at -20° C; stable for one year from the date of shipment. Non-hazardous. No MSDS required. Minimize repeated freezing and thawing.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.