



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# normal mouse IgG-HRP: sc-2748

## BACKGROUND

Santa Cruz Biotechnology offers a wide variety of control immunoglobulin and control sera for a large selection of species, including mouse, rabbit, goat, chicken, rat, hamster, canine, guinea pig and sheep. Control immunoglobulin and immunoglobulin conjugates are useful negative controls. Normal sera is offered to be used as blocking reagents. Santa Cruz Biotechnology offers affinity purified normal immunoglobulins and immunoglobulin conjugates for use as negative controls in applications including flow cytometry, immunohistochemistry, immunofluorescence, Western Blotting and immunoprecipitation. Agarose (AC) conjugated IgGs are provided for immunoprecipitation; horseradish peroxidase (HRP) conjugates are provided for Western Blotting and immunohistochemistry; and Biotin (B) conjugates are provided for immunohistochemistry. A broad range of fluorescent conjugated controls are also available for use in flow cytometry and immunofluorescence applications. Most control immunoglobulins are available as unconjugated controls or as FITC (fluorescein isothiocyanate), PE (phycoerythrin), PE-Cy5 (phycoerythrin-Cy5), PE-Cy7 (phycoerythrin-Cy7), APC (allophycocyanin) and APC-Cy7 (allophycocyanin-Cy7) conjugates. Additional conjugates include Alexa Fluor® 488, Alexa Fluor® 647, Alexa Fluor® 405, PerCP (peridinin chlorophyll protein complex) and PerCP-Cy5.5 (peridinin chlorophyll protein complex-Cy 5.5). Isotype specific control immunoglobulins include classes such as mouse IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>2b</sub>, IgG<sub>3</sub>, IgM and IgA, rat IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>2b</sub> and IgM, Armenian hamster IgG, and both goat and rabbit IgG.

## SOURCE

normal mouse IgG-HRP is an affinity purified, HRP (horseradish peroxidase) conjugated isotype control immunoglobulin from mouse.

## PRODUCT

Each vial contains 200 µg mouse IgG in PBS containing 40% glycerol, 1% protein stabilizer and < 0.01% thimerosal.

## APPLICATIONS

normal mouse IgG-HRP is recommended for use as an isotype control immunoglobulin in place of a target specific primary antibody of the same isotype (mouse IgG) by Western Blotting and immunohistochemical staining. To be used at an assay dependent dilution.

## RECOMMENDED SUPPORT PRODUCTS

- CrystalCruz® Cover Glasses, 22 x 50 mm, precleaned: sc-24975
- PBS (Phosphate Buffered Saline), powder, 1 packet: sc-24947
- Formaldehyde, 37% formaldehyde solution, 25 ml: sc-203049
- Hydrogen Peroxide, 30% solution, 100 ml: sc-203336
- Organo/Limonene Mount, non-toxic Permount alternative, 100 ml: sc-45087
- UltraCruz® Mounting Medium, aqueous-based, 10 ml: sc-24941
- ImmunoHistoMount, aqueous-based mounting medium, 30 ml: sc-45086
- Immuno In Situ Mount, for use with *in situ* hybridization, 30 ml: sc-45088
- Paraffin, for the preparation of tissue samples for staining, 500 g: sc-286633
- Xylenes, mixed isomers with ethylbenzene, 500 ml: sc-237422
- Hematoxylin, Gill's Formulation #2; nuclear counter stain, 100 ml: sc-24973

## STORAGE

Store at 4° C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## SELECT PRODUCT CITATIONS

1. Sattler, M. 1997. Steel factor induces tyrosine phosphorylation of CRKL and binding of CRKL to a complex containing c-Kit, phosphatidylinositol 3-kinase, and p120CBL. *J. Biol. Chem.* 272: 10248-10253.
2. Krones, A. 2001. Cross-talk between the signals hypoxia and glucose at the glucose response element of the L-type pyruvate kinase gene. *Endocrinology* 142: 2707-2718.
3. Schiller, M., et al. 2003. Cyclic adenosine 3',5'-monophosphate-elevating agents inhibit transforming growth factor—induced SMAD3/4-dependent transcription via a protein kinase A-dependent mechanism. *Oncogene.* 22: 8881-8890.
4. Onay-Besikci, A. 2004. gAd-globular head domain of adiponectin increases fatty acid oxidation in newborn rabbit hearts. *J. Biol. Chem.* 279: 44320-44326.
5. Fanara, P. 2004. *In vivo* measurement of microtubule dynamics using stable isotope labeling with heavy water. *J. Biol. Chem.* 279: 49940-49947.
6. Hashimoto, K., et al. 2007. Geranylgeranylacetone inhibits ovarian cancer progression *in vitro* and *in vivo*. *Biochem. Biophys. Res. Commun.* 356: 72-77.
7. Park, I.Y, et al. 2007. Aberrant epigenetic modifications in hepatocarcinogenesis induced by hepatitis B virus X protein. *Gastroenterology.* 132: 1476-1494.
8. Kedar, P.S., et al. 2008. Interaction between PARP-1 and ATR in mouse fibroblasts is blocked by PARP inhibition. *DNA Repair.* 7: 1787-1798.
8. Konduri, S.D., et al. 2009. Blockade of MGMT expression by O6 benzyl guanine leads to inhibition of pancreatic cancer growth and induction of apoptosis. *Clin. Cancer Res.* 15: 6087-6095.
9. Stella, R., et al. 2010. Cellular prion protein promotes regeneration of adult muscle tissue. *Mol. Cell. Biol.* 30: 4864-4876.
10. Tong, K.M., et al. 2011. Adiponectin increases MMP-3 expression in human chondrocytes through adipor1 signaling pathway. *J. Cell. Biochem.* 112: 1431-1440.
11. Choi, J.K., et al. 2011. Granulocyte macrophage-colony stimulating factor shows anti-apoptotic activity in neural progenitor cells via JAK/STAT5-Bcl-2 pathway. *Apoptosis.* 16: 127-134.
12. Muller, A.P., et al. 2012. Reduced brain insulin-like growth factor I function during aging. *Molecular and cellular neurosciences.* 49: 9-12.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

Alexa Fluor® is a registered trademark of Molecular Probes, Inc., Oregon, USA