

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

Sox-2 (h): 293T Lysate: sc-369948



BACKGROUND

Sox genes comprise a family of genes that are related to the mammalian sex determining gene SRY. These genes similarly contain sequences that encode for the HMG-box domain, which is responsible for the sequence-specific DNA-binding activity. Sox genes encode putative transcriptional regulators implicated in the decision of cell fates during development and the control of diverse developmental processes. The highly complex group of Sox genes cluster at a minimum of 40 different loci that rapidly diverged in various animal lineages. At present 30 Sox genes have been identified, and members of this family have been shown to be conserved during evolution and to play key roles during animal development. Some are involved in human diseases, including sex reversal.

REFERENCES

- Laudet, V., et al. 1993. Ancestry and diversity of the HMG box superfamily. Nucleic Acids Res. 21: 2493-2501.
- 2. Kuhlbrodt, K., et al. 1998. Sox-10, a novel transcriptional modulator in glial cells. J. Neurosci. 18: 237-250.
- 3. Arsic, N., et al. 1998. Characterisation and mapping of the human Sox-14 gene. Cytogenet. Cell Genet. 83: 139-146.
- 4. Osaki, E., et al. 1999. Identification of a novel Sry-related gene and its germ cell-specific expression. Nucleic Acids Res. 27: 2503-2510.
- 5. Sasai, Y. 2001. Roles of Sox factors in neural determination: conserved signaling in evolution? Int. J. Dev. Biol. 45: 321-326.

CHROMOSOMAL LOCATION

Genetic locus: SOX2 (human) mapping to 3q26.33.

PRODUCT

Sox-2 (h): 293T Lysate represents a lysate of human Sox-2 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

Sox-2 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive Sox-2 antibodies. Recommended use: 10-20 μl per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

Sox-2 (E-4) Alexa Fluor[®] 790: sc-365823 AF790 is recommended as a positive control antibody for Western Blot analysis of enhanced human Sox-2 expression in Sox-2 transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgGκ BP-HRP: sc-516102 or m-IgGκ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

DATA





Sox-2 (E-4): sc-365823. Western blot analysis of Sox-2 expression in non-transfected $({\bf A})$ and human Sox-2 transfected $({\bf B})$ 293T whole cell lysates.

Sox-2 (E-4): sc-365823. Western blot analysis of Sox-2 expression in non-transfected: sc-117752 (A) and human Sox-2 transfected: sc-369948 (B) 293T whole cell lysates.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.