



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



IFI-44 (h): 293T Lysate: sc-370929

BACKGROUND

The Interferon (IFN) family of proteins are able to alter the expression of a variety of target genes, thereby controlling a number of events within the cell. IFI-44 (interferon-induced protein 44), also known as p44 or MTAP44 (micro-tubule-associated protein 44), is a 444 amino acid protein that localizes to the cytoplasm and, upon induction by IFN- β s, aggregates to form microtubular structures. Human IFI-44 shares 97% sequence similarity with its chimp counterpart, suggesting a conserved role between species. The gene encoding IFI-44 maps to human chromosome 1, which spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

1. Takahashi, K., et al. 1990. Cloning, sequencing and expression in *Escherichia coli* of cDNA for a non-A, non-B hepatitis-associated microtubular aggregate protein. *J. Gen. Virol.* 71: 2005-2011.
2. Kitamura, A., et al. 1994. Induction of the human gene for p44, a hepatitis-C-associated microtubular aggregate protein, by interferon- α/β . *Eur. J. Biochem.* 224: 877-883.
3. Izmailova, E., et al. 2003. HIV-1 Tat reprograms immature dendritic cells to express chemoattractants for activated T cells and macrophages. *Nat. Med.* 9: 191-197.
4. Hwang, Y., et al. 2006. Genetic predisposition of responsiveness to therapy for chronic hepatitis C. *Pharmacogenomics* 7: 697-709.
5. Online Mendelian Inheritance in Man, OMIM™. 2006. Johns Hopkins University, Baltimore, MD. MIM Number: 610468. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Hallen, L.C., et al. 2007. Antiproliferative activity of the human IFN- α -inducible protein IFI44. *J. Interferon Cytokine Res.* 27: 675-680.
7. Mailaparambil, B., et al. 2008. Polymorphisms of interferons and their receptors in the genetics of severe RSV-associated diseases. *Arch. Virol.* 153: 2133-2137.

CHROMOSOMAL LOCATION

Genetic locus: IFI44 (human) mapping to 1p31.1.

PRODUCT

IFI-44 (h): 293T Lysate represents a lysate of human IFI-44 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

APPLICATIONS

IFI-44 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive IFI-44 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.