



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



elf3 ϵ (h2): 293T Lysate: sc-371433

BACKGROUND

The initiation of protein synthesis in eukaryotic cells is regulated by interactions between protein initiation factors and RNA molecules. Eukaryotic initiation factors (eIFs) are utilized in a sequence of reactions that lead to 80S ribosomal assembly and, ultimately, translation. The eukaryotic initiation factor-3 (eIF3) scaffolding structure is the largest of the eIF complexes and includes eIF3 α , eIF3 β , eIF3 γ , eIF3 δ , eIF3 ϵ , eIF3 ζ , eIF3 η and eIF3 θ , all of which function to control the assembly of the 40S ribosomal subunit. Association of eIF3 proteins with the 40S ribosomal subunit stabilizes eIF2-GTP-Met-tRNAiMet complex association and mRNA binding, and promotes dissociation of 80S ribosomes into 40S and 60S subunits, thereby promoting the assembly of the pre-initiation complex. Overexpression of eIF3 proteins is common in several cancers, suggesting a role for eIF3 proteins in tumorigenesis.

REFERENCES

- Valásek, L., Nielsen, K.H., Zhang, F., Fekete, C.A. and Hinnebusch, A.G. 2004. Interactions of eukaryotic translation initiation factor 3 (eIF3) subunit NIP1/c with eIF1 and eIF5 promote preinitiation complex assembly and regulate start codon selection. *Mol. Cell. Biol.* 24: 9437-9455.
- Peterson, T.R. and Sabatini, D.M. 2005. eIF3: a connecTOR of S6K1 to the translation preinitiation complex. *Mol. Cell* 20: 655-657.
- Dong, Z. and Zhang, J.T. 2006. Initiation factor eIF3 and regulation of mRNA translation, cell growth, and cancer. *Crit. Rev. Oncol. Hematol.* 59: 169-180.
- LeFebvre, A.K., Korneeva, N.L., Trutschl, M., Cvek, U., Duzan, R.D., Bradley, C.A., Hershey, J.W. and Rhoads, R.E. 2006. Translation initiation factor eIF4G-1 binds to eIF3 through the eIF3 ϵ subunit. *J. Biol. Chem.* 281: 22917-22932.
- Hinnebusch, A.G. 2006. eIF3: a versatile scaffold for translation initiation complexes. *Trends Biochem. Sci.* 31: 553-562.
- Masutani, M., Sonenberg, N., Yokoyama, S. and Imataka, H. 2007. Reconstitution reveals the functional core of mammalian eIF3. *EMBO J.* 26: 3373-3383.
- Zhang, L., Pan, X. and Hershey, J.W. 2007. Individual overexpression of five subunits of human translation initiation factor eIF3 promotes malignant transformation of immortal fibroblast cells. *J. Biol. Chem.* 282: 5790-5800.
- Sato, H., Masuda, M., Kanai, M., Tsukiyama-Kohara, K., Yoneda, M. and Kai, C. 2007. Measles virus N protein inhibits host translation by binding to eIF3-p40. *J. Virol.* 81: 11569-11576.
- Zhang, L., Smit-McBride, Z., Pan, X., Rheinhardt, J. and Hershey, J.W. 2008. An oncogenic role for the phosphorylated η -subunit of human translation initiation factor eIF3. *J. Biol. Chem.* 283: 24047-24060.

CHROMOSOMAL LOCATION

Genetic locus: EIF3F (human) mapping to 11p15.4.

PRODUCT

elf3 ϵ (h2): 293T Lysate represents a lysate of human eIF3 ϵ transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

elf3 ϵ (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive eIF3 ϵ antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.


eIF3 ϵ G-7: sc-390413 is recommended as a positive control antibody for Western Blot analysis of enhanced human eIF3 ϵ expression in eIF3 ϵ transfected 293T cells (starting dilution 1:100, dilution range 1:100-1:1,000).

RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:

1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048.

DATA



elf3 ϵ (G-7): sc-390413. Western blot analysis of eIF3 ϵ expression in non-transfected: sc-117752 (**A**) and human eIF3 ϵ transfected: sc-371433 (**B**) whole cell lysates.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.