



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TNF-IP 8 (h): 293T Lysate: sc-371558

BACKGROUND

TNF-IP 8, also known as TNFAIP8 (tumor necrosis factor α -induced protein 8), TNF-induced protein GG2-1, head and neck tumor and metastasis-related protein, NDED (NF κ B-inducible DED-containing protein) or SCC-S2, is a 198 amino acid cytoplasmic protein that exists as 3 alternatively spliced isoforms and belongs to the TNFAIP8 family. TNF-IP 8 acts as a negative mediator of apoptosis and may play a role in tumor progression. While it suppresses the TNF-mediated apoptosis by inhibiting caspase-8 activity, TNF-IP 8 does not suppress the processing of procaspase-8. TNF-IP 8 is expressed at high levels in spleen, lymph node, thymus, thyroid, bone marrow, placenta, and fetal liver, lung and kidney. The gene that encodes TNF-IP 8 maps to human chromosome 5q23.1.

REFERENCES

- Horrevoets, A.J., Fontijn, R.D., van Zonneveld, A.J., de Vries, C.J., ten Cate, J.W. and Pannekoek, H. 1999. Vascular endothelial genes that are responsive to tumor necrosis factor- α *in vitro* are expressed in atherosclerotic lesions, including inhibitor of apoptosis protein-1, stannin, and two novel genes. *Blood* 93: 3418-3431.
- Kumar, D., Whiteside, T.L. and Kasid, U. 2000. Identification of a novel tumor necrosis factor- α -inducible gene, SCC-S2, containing the consensus sequence of a death effector domain of fas-associated death domain-like interleukin-1 β -converting enzyme-inhibitory protein. *J. Biol. Chem.* 275: 2973-2978.
- You, Z., Ouyang, H., Lopatin, D., Polver, P.J. and Wang, C.Y. 2001. Nuclear factor- κ B-inducible death effector domain-containing protein suppresses tumor necrosis factor-mediated apoptosis by inhibiting caspase-8 activity. *J. Biol. Chem.* 276: 26398-26404.
- Kumar, D., Gokhale, P., Broustas, C., Chakravarty, D., Ahmad, I. and Kasid, U. 2004. Expression of SCC-S2, an antiapoptotic molecule, correlates with enhanced proliferation and tumorigenicity of MDA-MB 435 cells. *Oncogene* 23: 612-616.
- Online Mendelian Inheritance in Man, OMIM™. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 612111. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
- Zhang, L.J., Liu, X., Gafken, P.R., Kioussi, C. and Leid, M. 2009. A chicken ovalbumin upstream promoter transcription factor I (COUP-TFI) complex represses expression of the gene encoding tumor necrosis factor α -induced protein 8 (TNFAIP8). *J. Biol. Chem.* 284: 6156-6168.
- Zhang, S., Zhang, Y., Wei, X., Zhen, J., Wang, Z., Li, M., Miao, W., Ding, H., Du, P., Zhang, W., He, M. and Yi, F. 2010. Expression and regulation of a novel identified TNFAIP8 family is associated with diabetic nephropathy. *Biochim. Biophys. Acta* 1802: 1078-1086.
- Dong, Q.Z., Zhao, Y., Liu, Y., Wang, Y., Zhang, P.X., Jiang, G.Y., Dong, X.J., Cui, Q.Z. and Wang, E.H. 2010. Overexpression of SCC-S2 correlates with lymph node metastasis and poor prognosis in patients with non-small-cell lung cancer. *Cancer Sci.* 101: 1562-1569.

CHROMOSOMAL LOCATION

Genetic locus: TNFAIP8 (human) mapping to 5q23.1.

PRODUCT

TNF-IP 8 (h): 293T Lysate represents a lysate of human TNF-IP 8 transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

TNF-IP 8 (h): 293T Lysate is suitable as a Western Blotting positive control for human reactive TNF-IP 8 antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.