



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

OCML (h2): 293T Lysate: sc-372089

BACKGROUND

The family of EF-hand type Ca²⁺-binding proteins includes calbindin (previously designated vitamin D-dependent Ca²⁺-binding protein), S-100 α and β , calgranulins A (also designated MRP8), B (also designated MRP14) and C (S-100 like proteins) and the parvalbumin family members, including parvalbumin α and parvalbumin β , also designated oncomodulin (OCM). Structurally and evolutionarily conserved, parvalbumin α and OCM proteins are distinct in expression and function. Parvalbumin α , also designated parvalbumin (PV), is most abundantly expressed in fast-contracting muscles, with lower expression levels in brain and some endocrine tissues, including kidney and parathyroid. Research indicates that parvalbumin α plays a significant role in muscle relaxation. OCM was originally thought to have expression restricted to neoplastic tissues, early embryonic cells and certain tumor cell lines. Recent research shows that OCM is also expressed and secreted by macrophages where, in the retina, it binds to retinal ganglion cells (RGCs) and functions to promote axon regeneration. OCM has also been detected in the auditory sensory cells of the organ of Corti in mammals. In humans, two different loci on chromosome 7 have been identified as OCM and OCML. These genes encode proteins 109 amino acids in length which share 99% sequence identity.

REFERENCES

1. Pfyffer, G.E., Faivre-Bauman, A., Tixier-Vidal, A., Norman, A.W. and Heizmann, C.W. 1987. Developmental and functional studies of parvalbumin and calbindin D28K in hypothalamic neurons grown in serum-free medium. *J. Neurochem.* 49: 442-451.
2. Kagi, U., Chafouleas, J.G., Norman, A.W. and Heizmann, C.W. 1988. Developmental appearance of the Ca²⁺-binding proteins parvalbumin, calbindin D-28K, S-100 proteins and calmodulin during testicular development in the rat. *Cell Tissue Res.* 252: 359-365.
3. Muntener, M., Kaser, L., Weber, J. and Berchtold, M.W. 1995. Increase of skeletal muscle relaxation speed by direct injection of parvalbumin cDNA. *Proc. Natl. Acad. Sci. USA* 92: 6504-6508.
4. Pauls, T.L., Cox, J.A. and Berchtold, M.W. 1996. The Ca²⁺-binding proteins parvalbumin and oncomodulin and their genes: new structural and functional findings. *Biochim. Biophys. Acta* 1306: 39-54.
5. Cox, J.A., Durussel, I., Scott, D.J. and Berchtold, M.W. 1999. Remodeling of the AB site of rat parvalbumin and oncomodulin into a canonical EF-hand. *Eur. J. Biochem.* 264: 790-799.
6. Hackney, C.M., Mahendrasingam, S., Penn, A. and Fettiplace, R. 2005. The concentrations of calcium buffering proteins in mammalian cochlear hair cells. *J. Neurosci.* 25: 7867-7875.
7. Yin, Y., Henzl, M.T., Lorber, B., Nakazawa, T., Thomas, T.T., Jiang, F., Langer, R. and Benowitz, L.I. 2006. Oncomodulin is a macrophage-derived signal for axon regeneration in retinal ganglion cells. *Nat. Neurosci.* 9: 843-852.

CHROMOSOMAL LOCATION

Genetic locus: OCM2 (human) mapping to 7q21.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PRODUCT

OCML (h2): 293T Lysate represents a lysate of human OCML transfected 293T cells and is provided as 100 μ g protein in 200 μ l SDS-PAGE buffer.

APPLICATIONS

OCML (h2): 293T Lysate is suitable as a Western Blotting positive control for human reactive OCML antibodies. Recommended use: 10-20 μ l per lane.

Control 293T Lysate: sc-117752 is available as a Western Blotting negative control lysate derived from non-transfected 293T cells.

STORAGE

Store at -20° C. Repeated freezing and thawing should be minimized. Sample vial should be boiled once prior to use. Non-hazardous. No MSDS required.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.