



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MMP-10 siRNA (h): sc-41555

BACKGROUND

The matrix metalloproteinases (MMP) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, Fibronectin, Laminin and proteoglycan. Transcription of MMP genes is differentially activated by phorbol ester, lipopolysaccharide (LPS) or staphylococcal enterotoxin B (SEB). MMP catalysis requires both calcium and zinc. MMP-10, also known as Stromelysin-2, is expressed in small intestine and at lower levels in lung and heart. MMP-10 functions similarly to MMP-3 in that it can degrade Fibronectin and gelatins type I, III, IV and V, however its action on collagens III, IV and V is very weak. Significantly, expression of MMP-10 is upregulated in ras-transformed HaCaT II-4 keratinocytes, therefore enabling the cells to undergo epithelial-to-mesenchymal transition. This evidence suggests that MMP-10, as well as other matrix metalloproteinases, may play a significant role in tumor metastasis.

REFERENCES

1. Tsuda, T.T., et al. 1993. Isolation and characterization of a high molecular weight type IV collagenase isolated from human carcinoma tissue. *FEBS Lett.* 319: 35-39.
2. Online Mendelian Inheritance in Man, OMIM[™]. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 185260. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. McGowan, P.M., et al. 2008. Matrix metalloproteinase expression and outcome in patients with breast cancer: analysis of a published database. *Ann. Oncol.* 19: 1566-1572.
4. Järvinen, K., et al. 2008. Selective iNOS inhibitor 1400W enhances anti-catabolic IL-10 and reduces destructive MMP-10 in OA cartilage. Survey of the effects of 1400W on inflammatory mediators produced by OA cartilage as detected by protein antibody array. *Clin. Exp. Rheumatol.* 26: 275-282.
5. Páramo, J.A., et al. 2008. Association between matrix metalloproteinase-10 concentration and smoking in individuals without cardiovascular disease. *Rev. Esp. Cardiol.* 61: 1267-1273.
6. Eiseler, T., et al. 2009. Protein kinase D1 regulates matrix metalloproteinase expression and inhibits breast cancer cell invasion. *Breast Cancer Res.* 11: R13.

CHROMOSOMAL LOCATION

Genetic locus: MMP10 (human) mapping to 11q22.2.

PRODUCT

MMP-10 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MMP-10 shRNA Plasmid (h): sc-41555-SH and MMP-10 shRNA (h) Lentiviral Particles: sc-41555-V as alternate gene silencing products.

For independent verification of MMP-10 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41555A, sc-41555B and sc-41555C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MMP-10 siRNA (h) is recommended for the inhibition of MMP-10 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

MMP-10 (LA-12): sc-80197 is recommended as a control antibody for monitoring of MMP-10 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MMP-10 gene expression knockdown using RT-PCR Primer: MMP-10 (h)-PR: sc-41555-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Steele, M.M., et al. 2011. β 1 integrin regulates MMP-10 dependant tubulogenesis in human lymphatic endothelial cells. *Matrix Biol.* 30: 218-224.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.