



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MT-MMP-6 siRNA (m): sc-41574

BACKGROUND

The matrix metalloproteinases (MMPs) are a family of peptidase enzymes responsible for the degradation of extracellular matrix components, including collagen, gelatin, Fibronectin, Laminin and proteoglycan. MMP catalysis requires both calcium and zinc. Membrane-type matrix metalloproteinases are type I membrane proteins that function to activate other MMPs. MT-MMP activation appears to be mediated by members of the proprotein convertase family, suggesting that a proprotein convertase/MT-MMP/MMP cascade may be involved in the regulation of ECM turnover. MT-MMP-6 (also known as MMP-25, MT6-MMP or Leukolysin) is the second GPI-anchored proteinase in the MMP family. Similar to MMP-3, MT-MMP-6 is able to cleave type IV collagen, Fibronectin, fibrin and gelatin. However, MT-MMP-6 cannot cleave Laminin-1 or activate progelatinase B. MT-MMP 6 is expressed in leukocytes, lung and spleen.

REFERENCES

1. Pei, D. 1999. Leukolysin/MMP25/MT6-MMP: a novel matrix metalloproteinase specifically expressed in the leukocyte lineage. *Cell Res.* 9: 291-303.
2. Velasco, G., et al. 2000. Human MT6-matrix metalloproteinase: identification, progelatinase A activation, and expression in brain tumors. *Cancer Res.* 60: 877-882.
3. Kojima, S., et al. 2000. Membrane-type 6 matrix metalloproteinase (MT6-MMP, MMP-25) is the second glycosyl-phosphatidyl inositol (GPI)-anchored MMP. *FEBS Lett.* 480: 142-146.
4. English, W.R., et al. 2001. Catalytic activities of membrane-type 6 matrix metalloproteinase (MMP25). *FEBS Lett.* 491: 137-142.
5. Nie, J. and Pei, D. 2003. Direct activation of pro-matrix metalloproteinase-2 by leukolysin/membrane-type 6 matrix metalloproteinase/matrix metalloproteinase 25 at the Asn¹⁰⁹-Tyr bond. *Cancer Res.* 63: 6758-6762.
6. Online Mendelian Inheritance in Man, OMIM™. 2004. Johns Hopkins University, Baltimore, MD. MIM Number: 608482. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Sun, Q., et al. 2007. MMP25 (MT6-MMP) is highly expressed in human colon cancer, promotes tumor growth, and exhibits unique biochemical properties. *J. Biol. Chem.* 282: 21998-22010.
8. Sohail, A., et al. 2008. MT4-(MMP17) and MT6-MMP (MMP25), A unique set of membrane-anchored matrix metalloproteinases: properties and expression in cancer. *Cancer Metastasis Rev.* 27: 289-302.

CHROMOSOMAL LOCATION

Genetic locus: Mmp25 (mouse) mapping to 17 A3.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

MT-MMP-6 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MT-MMP-6 shRNA Plasmid (h): sc-41573-SH and MT-MMP-6 shRNA (h) Lentiviral Particles: sc-41573-V as alternate gene silencing products.

For independent verification of MT-MMP-6 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41573A, sc-41573B and sc-41573C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCL, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MT-MMP-6 siRNA (m) is recommended for the inhibition of MT-MMP-6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MT-MMP-6 gene expression knockdown using RT-PCR Primer: MT-MMP-6 (m)-PR: sc-41574-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.