



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# G $\alpha$ 12 siRNA (h): sc-41742

## BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (a photon, pheromone, odorant, hormone or neurotransmitter) while the effectors (i.e., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. In mammals, G protein  $\alpha$ ,  $\beta$  and  $\gamma$  polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their  $\alpha$  subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Four distinct classes of G $\alpha$  subunits have been identified; these include G $\alpha_s$ , G $\alpha_i$ , G $\alpha_q$  and G $\alpha_{12/13}$ . The two members of the fourth class of G $\alpha$  subunit proteins, G $\alpha_{12}$  and G $\alpha_{13}$ , are insensitive to ADP-ribosylation by pertussis toxin, share 67% identity with each other and less than 45% identity with other G $\alpha$  subunits and are widely expressed in a broad range of tissues.

## REFERENCES

1. Strathmann, M., et al. 1989. Diversity of the G protein family: sequences from five additional  $\alpha$  subunits in the mouse. *Proc. Natl. Acad. Sci. USA* 86: 7407-7409.
2. Strathmann, M.P., et al. 1991. G $\alpha_{12}$  and G $\alpha_{13}$  subunits define a fourth class of G protein  $\alpha$  subunits. *Proc. Natl. Acad. Sci. USA* 88: 5582-5586.
3. Simon, M.I., et al. 1991. Diversity of G proteins in signal transduction. *Science* 252: 802-808.
4. Cali, J.J., et al. 1992. Selective tissue distribution of G protein  $\gamma$  subunits, including a new form of the  $\gamma$  subunits identified by cDNA cloning. *J. Biol. Chem.* 267: 24023-24027.

## CHROMOSOMAL LOCATION

Genetic locus: GNA12 (human) mapping to 7p22.3.

## PRODUCT

G $\alpha_{12}$  siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see G $\alpha_{12}$  shRNA Plasmid (h): sc-41742-SH and G $\alpha_{12}$  shRNA (h) Lentiviral Particles: sc-41742-V as alternate gene silencing products.

For independent verification of G $\alpha_{12}$  (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41742A, sc-41742B and sc-41742C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

G $\alpha_{12}$  siRNA (h) is recommended for the inhibition of G $\alpha_{12}$  expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

G $\alpha_{12}$  (E-12): sc-515445 is recommended as a control antibody for monitoring of G $\alpha_{12}$  gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor G $\alpha_{12}$  gene expression knockdown using RT-PCR Primer: G $\alpha_{12}$  (h)-PR: sc-41742-PR (20  $\mu$ l, 509 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## SELECT PRODUCT CITATIONS

1. Huang, C., et al. 2011. Role of p115RhoGEF in the regulation of extracellular Ca<sup>2+</sup>-induced choline kinase activation and prostate cancer cell proliferation. *Int. J. Cancer* 128: 2833-2842.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.