



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# G<sub>α</sub> 15 siRNA (m): sc-41747

## BACKGROUND

Heterotrimeric G proteins function to relay information from cell surface receptors to intracellular effectors. Each of a very broad range of receptors specifically detects an extracellular stimulus (a photon, pheromone, odorant, hormone or neurotransmitter) while the effectors (i.e., adenylyl cyclase), which act to generate one or more intracellular messengers, are less numerous. In mammals, G protein  $\alpha$ ,  $\beta$  and  $\gamma$  polypeptides are encoded by at least 16, 4 and 7 genes, respectively. Most interest in G proteins has been focused on their  $\alpha$  subunits, since these proteins bind and hydrolyze GTP and most obviously regulate the activity of the best studied effectors. Four distinct classes of G<sub>α</sub> subunits have been identified; these include G<sub>s</sub>, G<sub>i</sub>, G<sub>q</sub> and G<sub>12/13</sub>. The G<sub>i</sub> class comprises all the known  $\alpha$  subunits that are susceptible to pertussis toxin modifications, including G<sub>αi-1</sub>, G<sub>αi-2</sub>, G<sub>αi-3</sub>, G<sub>αo</sub>, G<sub>αt1</sub>, G<sub>αt2</sub>, G<sub>αz</sub> and G<sub>αgust</sub>. Of these, the three G<sub>αi</sub> subtypes function to open atrial potassium channels. G<sub>α16</sub> is a member of the G<sub>q</sub> subfamily and is expressed specifically in hematopoietic cells.

## REFERENCES

1. Jones, D.T., et al. 1990. Biochemical characterization of three stimulatory GTP-binding proteins. *J. Biol. Chem.* 265: 2671-2676.
2. Amatruda, T.T., III., et al. 1991. G<sub>α16</sub>, a G protein  $\alpha$  subunit specifically expressed in hematopoietic cells. *Proc. Natl. Acad. Sci. USA* 88: 5587-5591.
3. Simon, M.I., et al. 1991. Diversity of G proteins in signal transduction. *Science* 252: 802-808.
4. McLaughlin, S.K., et al. 1992. Gustducin is a taste-cell-specific G protein closely related to the transducins. *Nature* 357: 563-569.
5. Cali, J.J., et al. 1992. Selective tissue distribution of G protein  $\gamma$  subunits, including a new form of the  $\gamma$  subunits identified by cDNA cloning. *J. Biol. Chem.* 267: 24023-24027.
6. von Weizsäcker, E., et al. 1992. Diversity among the  $\beta$  subunits of heterotrimeric GTP-binding proteins: characterization of a novel  $\beta$  subunit cDNA. *Biochem. Biophys. Res. Commun.* 183: 350-356.
7. Conklin, B.R. and Bourne, H.R. 1993. Structural elements of G<sub>α</sub> subunits that interact with G<sub>βγ</sub> receptors, and effectors. *Cell* 73: 631-641.

## CHROMOSOMAL LOCATION

Genetic locus: Gna15 (mouse) mapping to 10 C1.

## PRODUCT

G<sub>α</sub> 15 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see G<sub>α</sub> 15 shRNA Plasmid (m): sc-41747-SH and G<sub>α</sub> 15 shRNA (m) Lentiviral Particles: sc-41747-V as alternate gene silencing products.

For independent verification of G<sub>α</sub> 15 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41747A, sc-41747B and sc-41747C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

G<sub>α</sub> 15 siRNA (m) is recommended for the inhibition of G<sub>α</sub> 15 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor G<sub>α</sub> 15 gene expression knockdown using RT-PCR Primer: G<sub>α</sub> 15 (m)-PR: sc-41747-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.