



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7


mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



D5DR siRNA (m): sc-41935



The Power to Question

BACKGROUND

D5DR (dopamine receptor D5, DR1B, DRD1L2) is a G protein-coupled receptor of the D1-like family that stimulates adenyl cyclase and has a ten-fold higher affinity for dopamine than the D1 subtype. D5DR is present in neurons in the limbic regions of the brain. D5DR mRNA is most abundant in discrete cortical areas (layers II, IV and VI), the dentate gyrus and hippocampal subfields with low levels in the striatum. There are five major types of dopamine receptors. All are G protein-coupled metabotropic receptors and can be excitatory or inhibitory to the post-synaptic neuron. D1 (D1A, D1C, D1D) and D5 (D1B) receptors belong to the D1-like receptor family. The D2, D3 and D4 receptors belong to the D2-like receptor family. Activation of the D1-like family receptors couples to increases in cAMP and is typically excitatory, while D2-like activation reduces cAMP and is typically inhibitory. Significant age-related decline in dopamine receptor mRNAs occurs in the hippocampus and entorhinal cortex.

REFERENCES

- Beischlag, T.V., et al. 1995. The human dopamine D5 receptor gene: cloning and characterization of the 5'-flanking and promoter region. *Biochemistry* 34: 5960-5970.
- Nguyen-Legros, J., et al. 1999. Dopamine receptor localization in the mammalian retina. *Mol. Neurobiol.* 19: 181-204.
- Iwasiow, R.M., et al. 1999. Delineation of the structural basis for the activation properties of the dopamine D1 receptor subtypes. *J. Biol. Chem.* 274: 31882-31890.
- Luedtke, R.R., et al. 1999. Immunoblot and immunohistochemical comparison of murine monoclonal antibodies specific for the rat D1a and D1b dopamine receptor subtypes. *J. Neuroimmunol.* 101: 170-187.
- Montague, D.M., et al. 2001. Quantification of D1B(D5) receptors in dopamine D1A receptor-deficient mice. *Synapse* 39: 319-322.
- Lidow, M.S., et al. 2003. D1 dopamine receptors in the mouse prefrontal cortex: Immunocytochemical and cognitive neuropharmacological analyses. *Synapse* 47: 101-108.

CHROMOSOMAL LOCATION

Genetic locus: Drd5 (mouse) mapping to 5 B3.

PRODUCT

D5DR siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 µM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see D5DR shRNA Plasmid (m): sc-41935-SH and D5DR shRNA (m) Lentiviral Particles: sc-41935-V as alternate gene silencing products.

For independent verification of D5DR (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-41935A, sc-41935B and sc-41935C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 µl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 µl of RNase-free water makes a 10 µM solution in a 10 µM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

D5DR siRNA (m) is recommended for the inhibition of D5DR expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

D5DR (SG4-D1b): sc-33661 is recommended as a control antibody for monitoring of D5DR gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG_x BP-HRP: sc-516102 or m-IgG_x BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG_x BP-FITC: sc-516140 or m-IgG_x BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor D5DR gene expression knockdown using RT-PCR Primer: D5DR (m)-PR: sc-41935-PR (20 µl, 464 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.