



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Munc13-3 siRNA (h): sc-42024

BACKGROUND

Munc13 proteins (Munc13-1, Munc13-2, and Munc13-3) make up a family of highly homologous synaptic molecules that bind Syntaxin, an essential mediator of neurotransmitter release. Munc13 proteins contain phorbol ester binding C1- and C2-domains, which are regulatory domains for Ca²⁺, phospholipids and diacylglycerol. Munc13 proteins are primarily expressed by neurons, except for a ubiquitously expressed Munc13-2 splice variant. Munc13-1 is expressed by most neurons; it interacts with the N-terminal of Doc2 α , which is concentrated on the synaptic vesicle. Munc13-1 also interacts directly with msec7-1 to co-localize the two proteins at the active zone, a presynaptic, subcellular compartment with extremely high membrane turnover. Munc13-1 is essential for synaptic vesicle maturation and plays a role in the central priming function in synaptic vesicle exocytosis from glutamatergic synapses. Munc13-1 is concentrated in presynaptic terminals. Munc13-2 is expressed in rostral regions, whereas Munc13-3 is expressed primarily in the cerebellum.

REFERENCES

1. Brose, N., et al. 1995. Mammalian homologues of *Caenorhabditis elegans* unc-13 gene define novel family of C2-domain proteins. *J. Biol. Chem.* 270: 25273-25280.
2. Mochida, S., et al. 1998. Role of the Doc2 α -Munc13-1 interaction in the neurotransmitter release process. *Proc. Natl. Acad. Sci. USA* 95: 11418-11422.
3. Neeb, A., et al. 1999. Direct interaction between the ARF-specific guanine nucleotide exchange factor msec7-1 and presynaptic Munc13-1. *Eur. J. Cell Biol.* 78: 533-538.
4. Augustin, I., et al. 1999. Munc13-1 is essential for fusion competence of glutamatergic synaptic vesicles. *Nature* 400: 457-461.
5. Augustin, I., et al. 1999. Differential expression of two novel Munc13 proteins in rat brain. *Biochem. J.* 337: 363-371.
6. Koch, H., et al. 2000. Definition of Munc13-homology-domains and characterization of a novel ubiquitously expressed Munc13 isoform. *Biochem. J.* 349: 247-253.

CHROMOSOMAL LOCATION

Genetic locus: UNC13C (human) mapping to 15q21.3.

PRODUCT

Munc13-3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Munc13-3 shRNA Plasmid (h): sc-42024-SH and Munc13-3 shRNA (h) Lentiviral Particles: sc-42024-V as alternate gene silencing products.

For independent verification of Munc13-3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42024A, sc-42024B and sc-42024C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Munc13-3 siRNA (h) is recommended for the inhibition of Munc13-3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Munc13-3 gene expression knockdown using RT-PCR Primer: Munc13-3 (h)-PR: sc-42024-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.