



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Rtn-3 siRNA (h): sc-42219

## BACKGROUND

The Reticulon (Rtn) family consists of four members: Rtn-1 (also designated neuroendocrine-specific protein or NSP), Rtn-2 (also designated NSP-like-1), Rtn-3 (also designated NSP-like-2) and Nogo (also designated Rtn-4A). Reticulon proteins are anchored to the membranes of the endoplasmic reticulum through their common C-terminal regions. Localized on human chromosome 14q23.1, the gene encoding Rtn-1 is expressed as three isoforms: Rtn-1A (NSP-A), Rtn-1B (NSP-B) and Rtn-1C (NSP-C). The gene encoding human Rtn-2 is located on chromosome 19q13 and also encodes three isoforms. Rtn-2-A and Rtn-2-C are produced by the use of alternative initiation sites, whereas Rtn-2-B is an alternative splice variant of the Rtn-2-A isoform. Rtn-2-A and Rtn-2-B are highly expressed in brain, while Rtn-2-C is primarily expressed in skeletal muscle. The human Rtn-3 gene is located on chromosome 11q13.1 and is widely expressed, with the highest expression being in brain.

## REFERENCES

- Senden, N.H., et al. 1994. Subcellular localization and supramolecular organization of neuroendocrine-specific protein B (NSP-B) in small cell lung cancer. *Eur. J. Cell Biol.* 65: 341-353.
- van de Velde, H. J., et al. 1994. NSP-encoded reticulons are neuroendocrine markers of a novel category in human lung cancer diagnosis. *Cancer Res.* 54: 4769-4776.
- Geisler, J.G., et al. 1998. Molecular cloning of a novel mouse gene with predominant muscle and neural expression. *Mamm. Genome* 9: 274-282.
- Roebroek, A.J., et al. 1998. cDNA cloning, genomic organization, and expression of the human RTN2 gene, a member of a gene family encoding reticulons. *Genomics* 51: 98-106.
- Hens, J., et al. 1998. Neuronal differentiation is accompanied by NSP-C expression. *Cell Tissue Res.* 292: 229-237.
- Moreira, E.F., et al. 1999. Cloning of a novel member of the reticulon gene family (RTN3): gene structure and chromosomal localization to 11q13. *Genomics* 58: 73-81.

## CHROMOSOMAL LOCATION

Genetic locus: RTN3 (human) mapping to 11q13.1.

## PRODUCT

Rtn-3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rtn-3 shRNA Plasmid (h): sc-42219-SH and Rtn-3 shRNA (h) Lentiviral Particles: sc-42219-V as alternate gene silencing products.

For independent verification of Rtn-3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42219A, sc-42219B and sc-42219C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Rtn-3 siRNA (h) is recommended for the inhibition of Rtn-3 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

Rtn-3 (F-6): sc-374599 is recommended as a control antibody for monitoring of Rtn-3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker<sup>™</sup> Molecular Weight Standards: sc-2035, UltraCruz<sup>®</sup> Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rtn-3 gene expression knockdown using RT-PCR Primer: Rtn-3 (h)-PR: sc-42219-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.