



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

MiRP1 siRNA (m): sc-42510

BACKGROUND

Voltage-gated K⁺ channels in the plasma membrane control the repolarization and the frequency of action potentials in neurons, muscles, and other excitable cells. KCNE1 and KCNE2 (also known as MiRP1) are two single transmembrane domain β subunits of the delayed rectifier potassium channel IKr. In cardiac tissue, MiRP1 assembles with HERG, the pore-forming α subunit of IKr. In the brain, MiRP1 associates with KCNQ2 and accelerates the dissociation of KCNQ2 from the KCNQ2-KCNQ3 complex. MiRP1 also regulates the current amplitude and gating properties of the KCNQ1 K⁺ channel, and may assemble with KCNQ1 in the stomach to aid in K⁺ recycling, which is necessary for gastric acid secretion. The gene encoding human MiRP1 maps to chromosome 21q22.11. Missense mutations in the gene for MiRP1 result in congenital long QT syndrome and drug-induced cardiac arrhythmia.

REFERENCES

1. Kadenbach, B., et al. 1983. Separation of mammalian cytochrome c oxidase into 13 polypeptides by a sodium dodecyl sulfate-gel electrophoretic procedure. *Anal. Biochem.* 129: 517-521.
2. Shoffner, J.M. and Wallace, D.C. 1995. Oxidative Phosphorylation Diseases. In Scriver, C.R., Beaudet, A.L., Sly, W.S., Valle, D., eds., *The Metabolic and Molecular Basis of Inherited Disease*. New York: McGraw-Hill, 1535-1609.
3. Online Mendelian Inheritance in Man, OMIM[™]. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 516030. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
4. Cannino, G., et al. 2004. Analysis of cytochrome c oxidase subunits III and IV expression in developing rat brain. *Neuroscience* 128: 91-98.
5. Boerner, J.L., et al. 2004. Phosphorylation of Y845 on the epidermal growth factor receptor mediates binding to the mitochondrial protein cytochrome c oxidase subunit II. *Mol. Cell. Biol.* 24: 7059-7071.
6. Qian, J., et al. 2004. Role of the conserved arginine pair in proton and electron transfer in cytochrome c oxidase. *Biochemistry* 43: 5748-5756.
7. LocusLink Report (LocusID: 4512). <http://www.ncbi.nlm.nih.gov/LocusLink/>

CHROMOSOMAL LOCATION

Genetic locus: Kcne2 (mouse) mapping to 16 C4.

PRODUCT

MiRP1 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MiRP1 shRNA Plasmid (m): sc-42510-SH and MiRP1 shRNA (m) Lentiviral Particles: sc-42510-V as alternate gene silencing products.

For independent verification of MiRP1 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42510A, sc-42510B and sc-42510C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

MiRP1 siRNA (m) is recommended for the inhibition of MiRP1 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MiRP1 gene expression knockdown using RT-PCR Primer: MiRP1 (m)-PR: sc-42510-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.