



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

P2X4 siRNA (h): sc-42569

BACKGROUND

The P2X receptor family is comprised of ligand-gated ion channels that allow for the increased permeability of calcium into the cell in response to extracellular ATP. The seven P2X receptors, P2X1-P2X7, form either homomeric or heteromeric channels or both. They are characterized by intracellular amino- and carboxy-termini. P2X receptors are expressed in a wide variety of tissues, including neurons, prostate, bladder, pancreas, colon, testis and ovary. The major function of the P2X receptors is to mediate synaptic transmissions between neurons and to other tissues via the binding of extracellular ATP, which acts as a neurotransmitter. The P2X receptors may be involved in the onset of necrosis or apoptosis after prolonged exposure to high concentrations of extracellular ATP.

REFERENCES

1. Longhurst, P.A., et al. 1996. The human P_{2X1} receptor: molecular cloning, tissue distribution, and localization to chromosome 17. *Biochim. Biophys. Acta* 1308: 185-188.
2. Di Virgilio, F., et al. 1998. Cytolytic P2X purinoceptors. *Cell Death Differ.* 5: 191-199.
3. Alexander, K., et al. 1999. Allosteric modulation and accelerated resensitization of human P2X₃ receptors by cibacron blue. *J. Pharmacol. Exp. Ther.* 291: 1135-1142.
4. Burnstock, G. 2000. P2X receptors in sensory neurones. *Br. J. Anaesth.* 84: 476-488.
5. Oury, C., et al. 2000. A natural dominant negative P2X1 receptor due to deletion of a single amino acid residue. *J. Biol. Chem.* 275: 22611-22614.
6. Ding, S., et al. 2000. Inactivation of P2X₂ purinoceptors by divalent cations. *J. Physiol.* 522: 199-214.
7. North, R.A., et al. 2000. Pharmacology of cloned P2X receptors. *Annu. Rev. Pharmacol. Toxicol.* 40: 563-580.
8. Jabs, R., et al. 2000. Evidence for P2X₃, P2X₄, P2X₅ but not for P2X₇ containing purinergic receptors in Müller cells of the rat retina. *Brain Res. Mol. Brain Res.* 76: 205-210.

CHROMOSOMAL LOCATION

Genetic locus: P2RX4 (human) mapping to 12q24.31.

PRODUCT

P2X4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see P2X4 shRNA Plasmid (h): sc-42569-SH and P2X4 shRNA (h) Lentiviral Particles: sc-42569-V as alternate gene silencing products.

For independent verification of P2X4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42569A, sc-42569B and sc-42569C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

P2X4 siRNA (h) is recommended for the inhibition of P2X4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor P2X4 gene expression knockdown using RT-PCR Primer: P2X4 (h)-PR: sc-42569-PR (20 μ l, 428 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Freeman, K.W., et al. 2011. Regenerative protein thymosin β -4 is a novel regulator of purinergic signaling. *FASEB J.* 25: 907-915.
2. Chen, J.B., et al. 2012. Adenosine-5'-triphosphate up-regulates proliferation of human cardiac fibroblasts. *Br. J. Pharmacol.* 166: 1140-1150.
3. Zheng, K., et al. 2017. NOX2-mediated TFEB activation and vacuolization regulate lysosome-associated cell death induced by gypenoside L, a saponin isolated from *Gynostemma pentaphyllum*. *J. Agric. Food Chem.* 65: 6625-6637.
4. Chen, Y.J., et al. 2018. The suppressive effect of arsenic trioxide on TET2-FOXp3-Lyn-Akt axis-modulated MCL1 expression induces apoptosis in human leukemia cells. *Toxicol. Appl. Pharmacol.* 358: 43-55.
5. Ghalali, A., et al. 2020. PTEN and PHLPP crosstalk in cancer cells and in TGF β -activated stem cells. *Biomed. Pharmacother.* 127: 110112.

RESEARCH USE

For research use only, not for use in diagnostic procedures.