



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

KIR3.2 siRNA (h): sc-42618

BACKGROUND

G protein-coupled inwardly rectifying potassium channels (Kir3.1 through Kir3.4) are coupled to numerous neurotransmitter receptors in the brain and are abundantly expressed in the olfactory bulb, hippocampus, neocortex, dentate gyrus, cerebellar cortex and thalamus regions of the brain. Also known as GIRK, Kir3 potassium channels localize to the soma and dendrites as well as axons of neurons. Liberated $G_{\beta\gamma}$ subunits from G protein heterotrimers bind to and regulate Kir3 channel activity. $G_{\beta 3}$ - and $G_{\beta 4}$ -containing $G_{\beta\gamma}$ dimers bind directly to cytoplasmic domains of Kir3 proteins and increase the K^+ current while $G_{\beta 5}$ -containing $G_{\beta\gamma}$ dimers inhibit Kir3 K^+ current. Kir3 activity is also inhibited by tyrosine phosphorylation. Brain-derived neurotrophic factor activates receptor tyrosine kinase B, which then phosphorylates Kir3 tyrosine residues, effectively inactivating the Kir3 channels.

REFERENCES

1. Braun, A.P., et al. 1992. Activation of α 1-adrenoceptors modulates the inwardly rectifying potassium currents of mammalian atrial myocytes. *Pflugers Arch.* 421: 431-439.
2. Ponce, A., et al. 1996. G protein-gated inward rectifier K^+ channel proteins (GIRK1) are present in the soma and dendrites as well as in nerve terminals of specific neurons in the brain. *J. Neurosci.* 16: 1990-2001.
3. Farkas, R.H., et al. 1997. Neurotensin and dopamine D2 activation oppositely regulate the same K^+ conductance in rat midbrain dopaminergic neurons. *Neurosci. Lett.* 231: 21-24.
4. Lei, Q., et al. 2000. Activation and inhibition of G protein-coupled inwardly rectifying potassium (Kir3) channels by G protein by subunits. *Proc. Natl. Acad. Sci.* 97: 9771-9776.
5. Rogalski, S.L., et al. 2000. TrkB Activation by brain-derived neurotrophic factor inhibits the G protein-gated inward rectifier Kir3 by tyrosine phosphorylation of the channel. *J. Biol. Chem.* 275: 25082-25088.

CHROMOSOMAL LOCATION

Genetic locus: KCNJ6 (human) mapping to 21q22.13.

PRODUCT

KIR3.2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KIR3.2 shRNA Plasmid (h): sc-42618-SH and KIR3.2 shRNA (h) Lentiviral Particles: sc-42618-V as alternate gene silencing products.

For independent verification of KIR3.2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42618A, sc-42618B and sc-42618C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KIR3.2 siRNA (h) is recommended for the inhibition of KIR3.2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KIR3.2 gene expression knockdown using RT-PCR Primer: KIR3.2 (h)-PR: sc-42618-PR (20 μ l). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.