



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



NHE-4 siRNA (h): sc-42654

BACKGROUND

Na^+/H^+ exchangers-1–6 (Na^+/H^+ antiports, NHE-1–6) are integral membrane proteins that are expressed in most mammalian tissues, where they regulate intracellular pH and cell volume. NHEs mediate the secondary active extrusion of hydrogen (H^+) ions out of cells in exchange for extracellular sodium (Na^+). Excluding NHE-1, which is ubiquitously expressed, NHE isoforms 2–6 have distinct tissue- and cell type-dependent expression and inhibitory characteristics by amiloride analogs. Rat NHE-4 protein, also known as solute carrier family 9 isoform-4 and SLC9A4, is expressed in epithelial cells, kidney and stomach, and localizes to the basolateral membrane.

REFERENCES

1. Fliegel, L., et al. 1993. Cloning and analysis of the human myocardial Na^+/H^+ exchanger. *Mol. Cell. Biochem.* 125: 137-143.
2. Biemesderfer, D., et al. 1993. NHE-3: a Na^+/H^+ exchanger isoform of renal brush border. *Am. J. Physiol.* 265: 736-742.
3. Noel, J., et al. 1995. Hormonal regulation, pharmacology, and membrane sorting of vertebrate Na^+/H^+ exchanger isoforms. *Am. J. Physiol.* 268: 283-296.
4. Klanke, C.A., et al. 1995. Molecular cloning and physical and genetic mapping of a novel human Na^+/H^+ exchanger (NHE-5/SLC9A5) to chromosome 16q22.1. *Genomics* 25: 615-622.
5. Cox, G.A., et al. 1997. Sodium/hydrogen exchanger gene defect in slow-wave epilepsy mutant mice. *Cell* 91: 139-148.
6. Pizzonia, J.H., et al. 1998. Immunohistochemical characterization of Na^+/H^+ exchanger isoform NHE-4. *Am. J. Physiol.* 275: 510-517.

CHROMOSOMAL LOCATION

Genetic locus: SLC9A4 (human) mapping to 2q12.1.

PRODUCT

NHE-4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NHE-4 shRNA Plasmid (h): sc-42654-SH and NHE-4 shRNA (h) Lentiviral Particles: sc-42654-V as alternate gene silencing products.

For independent verification of NHE-4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42654A, sc-42654B and sc-42654C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NHE-4 siRNA (h) is recommended for the inhibition of NHE-4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NHE-4 gene expression knockdown using RT-PCR Primer: NHE-4 (h)-PR: sc-42654-PR (20 μl , 494 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.