



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



NHE-6 siRNA (m): sc-42659

BACKGROUND

Na^+/H^+ exchangers-1–6 (Na^+/H^+ antiports, NHE-1–6) are integral membrane proteins that are expressed in most mammalian tissues where they regulate intracellular pH and cell volume. NHE's mediate the secondary active extrusion of hydrogen (H^+) ions out of cells in exchange for extracellular sodium (Na^+). Excluding NHE-1, which is ubiquitously expressed, the NHE isoforms 2–6 have distinct tissue- and cell type-dependent expression, and inhibitory characteristics by amiloride analogs.

REFERENCES

- Fliegel, L., et al. 1993. Cloning and analysis of the human myocardial Na^+/H^+ exchanger. *Mol. Cell. Biochem.* 125: 137-143.
- Biemesderfer, D., et al. 1993. NHE3: a Na^+/H^+ exchanger isoform of renal brush border. *Am. J. Physiol.* 265: 736-742.
- Klanke, C.A., et al. 1995. Molecular cloning and physical and genetic mapping of a novel human Na^+/H^+ exchanger (NHE5/SLC9A5) to chromosome 16q22.1. *Genomics* 25: 615-622.
- Noel, J., et al. 1995. Hormonal regulation, pharmacology, and membrane sorting of vertebrate Na^+/H^+ exchanger isoforms. *Am. J. Physiol.* 268: 283-296.
- Cox, G.A., et al. 1997. Sodium/hydrogen exchanger gene defect in slow-wave epilepsy mutant mice. *Cell* 91: 139-148.
- Baird, N.R., et al. 1999. Molecular cloning, genomic organization, and functional expression of Na^+/H^+ exchanger isoform 5 (NHE5) from human brain. *J. Biol. Chem.* 274: 4377-4382.

CHROMOSOMAL LOCATION

Genetic locus: Slc9a6 (mouse) mapping to X A5.

PRODUCT

NHE-6 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NHE-6 shRNA Plasmid (m): sc-42659-SH and NHE-6 shRNA (m) Lentiviral Particles: sc-42659-V as alternate gene silencing products.

For independent verification of NHE-6 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42659A, sc-42659B and sc-42659C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NHE-6 siRNA (m) is recommended for the inhibition of NHE-6 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NHE-6 gene expression knockdown using RT-PCR Primer: NHE-6 (m)-PR: sc-42659-PR (20 μl , 526 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.