



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



TRPC1 siRNA (h): sc-42664



The Power to Question

BACKGROUND

Transient receptor potential cation (TRPC) channels are a superfamily of six transmembrane segment-spanning, gated cation channels. TRPC subtypes mediate store-operated Ca^{2+} entry, a process involving Ca^{2+} influx and replenishment of Ca^{2+} stores formerly emptied through the action of inositol 1,4,5-trisphosphate production and other Ca^{2+} mobilizing agents. TRPC ion channels influence calcium-depletion induced calcium influx processes in response to chemo-, mechano- and osmoregulatory events. Human TRPC1 protein is a 793 amino acid cation channel that is expressed in fetal and adult brain, and adult heart, testis and ovary, where it may influence store-operated Ca^{2+} entry as a component of capacitative calcium entry (CCE) complexes. The activation of store-mediated Ca^{2+} entry in human cells occurs through the association between inositol 1,4,5-trisphosphate receptors and TRPC1.

CHROMOSOMAL LOCATION

Genetic locus: TRPC1 (human) mapping to 3q23.

PRODUCT

TRPC1 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfactions. Also see TRPC1 shRNA Plasmid (h): sc-42664-SH and TRPC1 shRNA (h) Lentiviral Particles: sc-42664-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TRPC1 siRNA (h) is recommended for the inhibition of TRPC1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

GENE EXPRESSION MONITORING

TRPC1 (E-6): sc-133076 is recommended as a control antibody for monitoring of TRPC1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRPC1 gene expression knockdown using RT-PCR Primer: TRPC1 (h)-PR: sc-42664-PR (20 μl , 491 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Bair, A.M., et al. 2009. Ca^{2+} entry via TRPC channels is necessary for thrombin-induced NF κ B activation in endothelial cells through AMP-activated protein kinase and protein kinase C δ . *J. Biol. Chem.* 284: 563-574.
- Thippegowda, P.B., et al. 2010. Ca^{2+} influx via TRPC channels induces NF κ B-dependent A20 expression to prevent thrombin-induced apoptosis in endothelial cells. *Am. J. Physiol., Cell Physiol.* 298: C656-C664.
- Sobradillo, D., et al. 2014. A reciprocal shift in transient receptor potential channel 1 (TRPC1) and stromal interaction molecule 2 (STIM2) contributes to Ca^{2+} remodeling and cancer hallmarks in colorectal carcinoma cells. *J. Biol. Chem.* 289: 28765-28782.
- Bodiga, V.L., et al. 2016. Intracellular zinc status influences cisplatin-induced endothelial permeability through modulation of PKC α , NF κ B and ICAM-1 expression. *Eur. J. Pharmacol.* 791: 355-368.
- Guéguinou, M., et al. 2016. SK3/TRPC1/Orai1 complex regulates SOCE-dependent colon cancer cell migration: a novel opportunity to modulate anti-EGFR mAb action by the alkyl-lipid Ohmeline. *Oncotarget* 7: 36168-36184.
- Wang, Y., et al. 2016. TRPC1/TRPC3 channels mediate lysophosphatidyl-choline-induced apoptosis in cultured human coronary artery smooth muscles cells. *Oncotarget* 7: 50937-50951.
- He, D., et al. 2020. TRPC1 participates in the HSV-1 infection process by facilitating viral entry. *Sci. Adv.* 6: eaaz3367.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.