



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# TRCP4 siRNA (m): sc-42669

## BACKGROUND

Transient receptor potential cation (TRPC) channels are a superfamily of six transmembrane segment-spanning, gated cation channels. TRPC subtypes mediate store-operated  $Ca^{2+}$  entry, a process involving  $Ca^{2+}$  influx and replenishment of  $Ca^{2+}$  stores formerly emptied through the action of inositol 1,4,5-trisphosphate production and other  $Ca^{2+}$  mobilizing agents. TRPC channels influence calcium-depletion induced calcium influx processes in response to chemo-, mechano- and osmoregulatory events. Human TRPC4 protein, also known as Trp4, functions as a cation channel and is a constituent of native store-operated  $Ca^{2+}$ -permeable channels. In the presence of elevated  $Ca^{2+}$  concentrations, TRPC4 binds Calmodulin (CaM) at an interface which comprises amino acids 688-759 and 786-848 of TRPC4. The ability of TRPC4 to increase inwardly rectifying  $K^+$  currents suggests that TRPC4 may contribute to the formation of a novel  $K^+$  channel or upregulate endogenous inwardly rectifying  $K^+$  channel expression or activity.

## REFERENCES

1. Philipp, S., et al. 1998. A novel capacitative calcium entry channel expressed in excitable cells. *EMBO J.* 17: 4274-4282.
2. Harteneck, C., et al. 2000. From worm to man: three subfamilies of TRP channels. *Trends Neurosci.* 23: 159-166.
3. Hofmann, T., et al. 2000. Transient receptor potential channels as molecular substrates of receptor-mediated cation entry. *J. Mol. Med.* 78: 14-25.
4. McKay, R.R., et al. 2000. Cloning and expression of the human transient receptor potential 4 (Trp4) gene: localization and functional expression of human Trp4 and Trp3. *Biochem. J.* 351: 735-746.
5. Zhang, Z., et al. 2001. Increased inwardly rectifying potassium currents in HEK-293 cells expressing murine transient receptor potential 4. *Biochem. J.* 354: 717-725.
6. Trost, C., et al. 2001. The transient receptor potential, Trp4, cation channel is a novel member of the family of calmodulin binding proteins. *Biochem. J.* 355: 663-670.

## CHROMOSOMAL LOCATION

Genetic locus: *Trpc4* (mouse) mapping to 3 C.

## PRODUCT

TRPC4 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRPC4 shRNA Plasmid (m): sc-42669-SH and TRPC4 shRNA (m) Lentiviral Particles: sc-42669-V as alternate gene silencing products.

For independent verification of TRPC4 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42669A, sc-42669B and sc-42669C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}$  C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}$  C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

TRPC4 siRNA (m) is recommended for the inhibition of TRPC4 expression in mouse cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRPC4 gene expression knockdown using RT-PCR Primer: TRPC4 (m)-PR: sc-42669-PR (20  $\mu$ l, 521 bp). Annealing temperature for the primers should be  $55-60^{\circ}$  C and the extension temperature should be  $68-72^{\circ}$  C.

## SELECT PRODUCT CITATIONS

1. Jeon, J.P., et al. 2013. Activation of TRPC4 $\beta$  by  $G_{\alpha_i}$  subunit increases  $Ca^{2+}$  selectivity and controls neurite morphogenesis in cultured hippocampal neuron. *Cell Calcium* 54: 307-319.
2. Chen, C., et al. 2017. Critical role of TRPC1 in thyroid hormone-dependent dopaminergic neuron development. *Biochim. Biophys. Acta Mol. Cell Res.* 1864: 1900-1912.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.