



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



CD53 siRNA (h): sc-42796



BACKGROUND

The CD53 antigen is a member of the tetraspanin membrane protein family that is expressed in the lymphoid-myeloid lineage. The tetraspanin superfamily (CD9, CD37, CD53, CD63, CD81 and CD82) comprises a group of cell-surface proteins that are involved in cell activation and signal transduction as well as in cell adhesion, motility and metastasis. Tetraspanin transmembrane proteins have a metastasis suppressor effect by acting as cell motility brakes in tumor cells. Human neutrophils express high levels of CD53, an N-glycosylated pan-leukocyte antigen and the true homolog of the rat MRC-OX-44 antigen. CD53 is expressed on B cells, monocytes, macrophages, neutrophils, single (CD4 or CD8) positive thymocytes and peripheral T cells.

REFERENCES

- Bazil, V., Stefanová, I., Hilgert, I., Kristofová, H., Vanek, S., Bukovský, A. and Horejsí, V. 1989. Monoclonal antibodies against human leucocyte antigens. III. Antibodies against CD45R, CD6, CD44 and two newly described broadly expressed glycoproteins MEM-53 and MEM-102. *Folia Biol.* 35: 289-297.
- Angelisova, P., Vlcek, C., Stefanova, I., Lipoldova, M. and Horejsi, V. 1990. The human leucocyte surface antigen CD53 is a protein structurally similar to the CD37 and MRC-OX-44 antigens. *Immunogenetics* 32: 281-285.
- Olweus, J., Lund-Johansen, F. and Horejsi, V. 1993. CD53, a protein with four membrane-spanning domains, mediates signal transduction in human monocytes and B cells. *J. Immunol.* 151: 707-716.
- Rasmussen, A.M., Blomhoff, H.K., Stokke, T., Horejsi, V. and Smeland, E.B. 1994. Cross-linking of CD53 promotes activation of resting human B lymphocytes. *J. Immunol.* 153: 4997-5007.
- Okochi, H., Kato, M., Nashiro, K., Yoshie, O., Miyazono, K. and Furue, M. 1997. Expression of tetra-spans transmembrane family (CD9, CD37, CD53, CD63, CD81 and CD82) in normal and neoplastic human keratinocytes: an association of CD9 with $\alpha\beta 1$ integrin. *Br. J. Dermatol.* 137: 856-863.

CHROMOSOMAL LOCATION

Genetic locus: CD53 (human) mapping to 1p13.3.

PRODUCT

CD53 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfactions. Also see CD53 shRNA Plasmid (h): sc-42796-SH and CD53 shRNA (h) Lentiviral Particles: sc-42796-V as alternate gene silencing products.

For independent verification of CD53 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42796A, sc-42796B and sc-42796C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CD53 siRNA (h) is recommended for the inhibition of CD53 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

CD53 (B-12): sc-390185 is recommended as a control antibody for monitoring of CD53 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:
1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CD53 gene expression knockdown using RT-PCR Primer: CD53 (h)-PR: sc-42796-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.