



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# DC-SIGNR siRNA (h): sc-42859

## BACKGROUND

Dendritic cells (DC) are antigen-presenting immune system cells that are present on peripheral mucosal tissues and migrate to lymphoid tissues. DC-SIGN (DC-specific ICAM-3 grabbing nonintegrin) is a type II membrane protein that is exclusively expressed by DC. DC-SIGN, also designated CD209, binds to ICAM-3 to mediate the initial interaction between DC and resting T cells through the immunological synapse. The DC that are present in the initial sites of HIV-1 infection capture HIV-1 through DC-SIGN, which then facilitates the migration of DC to areas of T cell-rich secondary lymphoid organs, where it promotes efficient *trans* HIV-1 infection of these T cells. DC-SIGNR (DC-SIGN-related molecule), also designated CD209L and L-SIGN (liver/lymph node-specific ICAM-3 grabbing nonintegrin), is a type II integral membrane protein that is 77% identical to DC-SIGN. It is expressed on sinusoidal endothelial cells and binds the E2 glycoproteins of the hepatitis C virus.

## REFERENCES

1. Fauci, A. 1996. Host factors and the pathogenesis of HIV-induced disease. *Nature* 384: 529-534.
2. Yokoyama-Kobayashi, M., et al. 1999. Selection of cDNAs encoding putative type II membrane proteins on the cell surface from a human full-length cDNA bank. *Gene* 228: 161-167.
3. Soilleux, E.J., et al. 2000. DC-SIGN; a related gene, DC-SIGNR; and CD23 form a cluster on 19p13. *J. Immunol.* 165: 2937-2942.
4. Bashirova, A.A., et al. 2001. A dendritic cell-specific intercellular adhesion molecule 3-grabbing nonintegrin (DC-SIGN)-related protein is highly expressed on human liver sinusoidal endothelial cells and promotes HIV-1 infection. *J. Exp. Med.* 193: 671-678.
5. Pöhlmann, S., et al. 2001. DC-SIGNR, a DC-SIGN homolog expressed in endothelial cells, binds to human and simian immunodeficiency viruses and activates infection in *trans*. *Proc. Natl. Acad. Sci. USA* 98: 2670-2675.
6. Lai, W.K., et al. 2006. Expression of DC-SIGN and DC-SIGNR on human sinusoidal endothelium: a role for capturing hepatitis C virus particles. *Am. J. Pathol.* 169:200-208.

## CHROMOSOMAL LOCATION

Genetic locus: CLEC4M (human) mapping to 19p13.2.

## PRODUCT

DC-SIGNR siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DC-SIGNR shRNA Plasmid (h): sc-42859-SH and DC-SIGNR shRNA (h) Lentiviral Particles: sc-42859-V as alternate gene silencing products.

For independent verification of DC-SIGNR (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42859A, sc-42859B and sc-42859C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

DC-SIGNR siRNA (h) is recommended for the inhibition of DC-SIGNR expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## GENE EXPRESSION MONITORING

DC-SIGNR (C-10): sc-271086 is recommended as a control antibody for monitoring of DC-SIGNR gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG $\kappa$  BP-HRP: sc-516102 or m-IgG $\kappa$  BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DC-SIGNR gene expression knockdown using RT-PCR Primer: DC-SIGNR (h)-PR: sc-42859-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.