



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

EAT-2 siRNA (h): sc-42970

BACKGROUND

The pathogenesis of the Ewing sarcoma family of tumors is characterized by the presence of an EWS/FLI1 fusion gene following a translocation between chromosomes 11 and 22, which results in the expression of a chimeric protein. Originally isolated from Ewing's sarcoma tumor cells lines, the EWS/FLI1 activated transcript 2 (EAT-2) protein is an intracellular signaling protein that is expressed in immune cells, including macrophages and B lymphocytes. EAT-2 is expressed in NIH/3T3 cells within 4-8 hours of EWS/FLI1 induction, suggesting a potential role for EAT-2 in the oncogenesis of Ewing's sarcoma. EAT-2 binds members of the signaling lymphocytic-activation molecule (SLAM) family of immune receptors, which are present in varying levels in immune cells. Specifically, EAT-2 plays a role in controlling the signal transduction of antigen-presenting cells by binding to SLAM family members CD150, CD244, CD84 and CD229, which contain conserved tyrosine motifs in their cytoplasmic tails.

REFERENCES

1. Thompson, A.D., Braun, B.S., Arvand, A., Stewart, S.D., May, W.A., Chen, E., Korenberg, J. and Denny, C. 1996. EAT-2 is a novel SH2 domain containing protein that is up regulated by Ewing's sarcoma EWS/FLI1 fusion gene. *Oncogene* 13: 2649-2658.
2. West, D.C. 2000. Ewing sarcoma family of tumors. *Curr. Opin. Oncol.* 12: 323-329.
3. Morra, M., Lu, J., Poy, F., Martin, M., Sayos, J., Calpe, S., Gullo, C., Howie, D., Rietdijk, S., Thompson, A., Coyle, A.J., Denny, C., Yaffe, M.B., Engel, P., Eck, M.J. and Terhorst, C. 2001. Structural basis for the interaction of the free SH2 domain EAT-2 with SLAM receptors in hematopoietic cells. *EMBO J.* 20: 5840-5852.
4. Veillette, A. 2002. The SAP family: a new class of adaptor-like molecules that regulates immune cell functions. *Sci. STKE* 120: 8.
5. Fraser, C.C., Howie, D., Morra, M., Qiu, Y., Murphy, C., Shen, Q., Gutierrez-Ramos, J.C., Coyle, A., Kingsbury, G.A. and Terhorst, C. 2002. Identification and characterization of SF2000 and SF2001, two new members of the immune receptor SLAM/CD2 family. *Immunogenetics* 53: 843-850.

CHROMOSOMAL LOCATION

Genetic locus: SH2D1B (human) mapping to 1q23.3.

PRODUCT

EAT-2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see EAT-2 shRNA Plasmid (h): sc-42970-SH and EAT-2 shRNA (h) Lentiviral Particles: sc-42970-V as alternate gene silencing products.

For independent verification of EAT-2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42970A, sc-42970B and sc-42970C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

EAT-2 siRNA (h) is recommended for the inhibition of EAT-2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor EAT-2 gene expression knockdown using RT-PCR Primer: EAT-2 (h)-PR: sc-42970-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Enose-Akahata, Y., Matsuura, E., Oh, U. and Jacobson, S. 2009. High expression of CD244 and SAP regulated CD8 T cell responses of patients with HTLV-I associated neurologic disease. *PLoS Pathog.* 5: e1000682.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.