



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

TCL-1A siRNA (h): sc-42988

BACKGROUND

T cell leukemia/lymphoma protein 1A (TCL-1A), also known as p14^{TCL-1}, is a product of the TCL-1 gene that is involved in T cell prolymphocytic leukemia (T-PLL). T-PLL is a rare form of mature T cell leukemia, which is consistently associated with chromosomal rearrangements characterized by the juxtaposition of the TCRA locus on chromosome 14q11 and the TCL1A gene on 14q32.13. TCL-1A is a member of a unique family of β -barrel proteins that bind small hydrophobic ligands and function in cell regulation. TCL-1A is an all- β protein containing an eight-stranded antiparallel β -barrel which consists of two four-stranded β -meander motifs. The two motifs are related by a twofold axis and connected by a long loop. TCL-1A forms a tight crystallographic dimer. TCL-1A is expressed in pre-B cells, in immature thymocytes, at low levels in activated T cells and in the cytoplasm.

REFERENCES

- Gritti, C., et al. 1998. Transgenic mice for MTCP-1 develop T cell prolymphocytic leukemia. *Blood* 92: 368-373.
- Fu, Z.Q., et al. 1998. Crystal structure of MTCP-1: implications for role of TCL-1 and MTCP-1 in T cell malignancies. *Proc. Natl. Acad. Sci. USA* 95: 3413-3418.
- Hoh, F., et al. 1998. Crystal structure of p14^{TCL-1}, an oncogene product involved in T cell prolymphocytic leukemia, reveals a novel β -barrel topology. *Structure* 6: 147-155.
- Du Bois, G.C., et al. 1998. Purification and characterization of recombinant forms of TCL-1 and MTCP-1 proteins. *Protein Expr. Purif.* 12: 215-225.

CHROMOSOMAL LOCATION

Genetic locus: TCL1A (human) mapping to 14q32.13.

PRODUCT

TCL-1A siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCL-1A shRNA Plasmid (h): sc-42988-SH and TCL-1A shRNA (h) Lentiviral Particles: sc-42988-V as alternate gene silencing products.

For independent verification of TCL-1A (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-42988A, sc-42988B and sc-42988C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TCL-1A siRNA (h) is recommended for the inhibition of TCL-1A expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

TCL-1A (A-6): sc-393436 is recommended as a control antibody for monitoring of TCL-1A gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCL-1A gene expression knockdown using RT-PCR Primer: TCL-1A (h)-PR: sc-42988-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- Motiwala, T., et al. 2011. AP-1 elements and TCL1 protein regulate expression of the gene encoding protein tyrosine phosphatase PTPROT in leukemia. *Blood* 118: 6132-6140.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.