



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## TCL-1B3a siRNA (m): sc-42994

### BACKGROUND

T cell leukemia/lymphoma protein 1B, TCL-1B, (also designated TML1, TCL/MTCP1-like 1, syncytiotrophoblast-specific protein and SYN-1), is a protein involved in T cell prolymphocytic leukemia (T-PLL). TCL-1B is located within the region on chromosome 14q32.1 which is frequently involved in chromosomal translocations and inversions with one of the T cell receptor loci in human T cell leukemias and lymphomas. TCL-1B is activated by chromosomal rearrangements involving the TCL1 locus. In mouse, TCL-1B is represented by five homologues, TCL-1B1-5. The crystal structure of TCL-1B suggests that it may play a role in the transport of small molecules such as retinoids, nucleotides and fatty acids. TCL-1B is found in both the nucleus and the cytoplasm of normal bone marrow and peripheral lymphocytes.

### PRODUCT

TCL-1B3a siRNA (m) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCL-1B3a shRNA Plasmid (m): sc-42994-SH and TCL-1B3a shRNA (m) Lentiviral Particles: sc-42994-V as alternate gene silencing products.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}$  C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}$  C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

TCL-1B3a siRNA (m) is recommended for the inhibition of TCL-1B3a expression in mouse cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCL-1B3a gene expression knockdown using RT-PCR Primer: TCL-1B3a (m)-PR: sc-42994-PR (20  $\mu$ l). Annealing temperature for the primers should be  $55-60^{\circ}$  C and the extension temperature should be  $68-72^{\circ}$  C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.