



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ICAM-5 siRNA (m): sc-43035

BACKGROUND

Cell adhesion molecules are a family of closely related cell surface glycoproteins involved in cell-cell interactions during growth. They are thought to play important, yet separate, roles in embryogenesis and development. The intracellular adhesion molecule-1 (ICAM-1), also referred to as CD54, is an integral membrane protein of the immunoglobulin superfamily and recognizes the α 1/ β 2 and α M/ β 2 integrins. ICAM-2 functions as a ligand for lymphocyte function-associated antigen-1 (LFA-1) and is involved in leukocyte adhesion. ICAM-3 is highly expressed on the surface of human eosinophils and, when bound to ligand, may inhibit eosinophil inflammatory responses and survival. ICAM-4, also known as LW glycoprotein, interacts with the integrins α L/ β 2, α M/ β 2, α 4/ β 1, the α V family and α IIb/ β 3. Selective binding to different integrins may be pathologically relevant in a number of red blood cell-associated diseases. Lastly, ICAM-5, expressed on telencephalic neurons, binds CD11a/CD18 and thus may act as an adhesion molecule for leukocyte binding in the central nervous system.

REFERENCES

- Jorgensen, O.S. 1995. Neural cell adhesion molecule (NCAM) as a quantitative marker in synaptic remodeling. *Neurochem. Res.* 20: 533-547.
- Edelman, G.M. and Jones, F.S. 1995. Developmental control of NCAM expression by HOX and PAX gene products. *Philos. Trans. R. Soc. Lond., B, Biol. Sci.* 349: 305-312.
- Briskin, M.J., Rott, L. and Butcher, E.C. 1996. Structural requirements for mucosal vascular addressin binding to its lymphocyte receptor α 4/ β 7. Common themes among integrin-Ig family interactions. *J. Immunol.* 156: 719-726.
- Heiska, L., Kantor, C., Parr, T., Critchley, D.R., Vilja, P., Gahmberg, C.G. and Carpen, O. 1996. Binding of the cytoplasmic domain of intercellular adhesion molecule-2 (ICAM-2) to α -actinin. *J. Biol. Chem.* 271: 26214-26219.
- Tian, L., Kilgannon, P., Yoshihara, Y., Mori, K., Gallatin, W.M., Carpen, O. and Gahmberg, C.G. 2000. Binding of T lymphocytes to hippocampal neurons through ICAM-5 (telencephalin) and characterization of its interaction with the leukocyte integrin CD11a/CD18. *Eur. J. Immunol.* 30: 810-818.
- Kessel JM, Gern, J.E., Vrtis, R.F., Sedgwick, J.B. and Busse, W.W. 2003. Ligation of intercellular adhesion molecule 3 inhibits GM-CSF production by human eosinophils. *J. Allergy Clin. Immunol.* 111: 1024-1031.
- Mankelov, T.J., Spring, F.A., Parsons, S.F., Brady, R.L., Mohandas, N., Chasis, J.A. and Anstee, D.J. 2003. Identification of critical amino-acid residues on the erythroid intercellular adhesion molecule-4 (ICAM-4) mediating adhesion to α V integrins. *Blood* 2003 103: 1503-1508.

CHROMOSOMAL LOCATION

Genetic locus: *Icam5* (mouse) mapping to 9 A3.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ICAM-5 siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ICAM-5 shRNA Plasmid (m): sc-43035-SH and ICAM-5 shRNA (m) Lentiviral Particles: sc-43035-V as alternate gene silencing products.

For independent verification of ICAM-5 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43035A, sc-43035B and sc-43035C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ICAM-5 siRNA (m) is recommended for the inhibition of ICAM-5 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ICAM-5 gene expression knockdown using RT-PCR Primer: ICAM-5 (m)-PR: sc-43035-PR (20 μ l, 428 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.