



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

α -actinin-4 siRNA (h): sc-43101

BACKGROUND

The spectrin gene family encodes a diverse group of cytoskeletal proteins that include spectrins, dystrophins and α -actinins. There are four tissue-specific α -actinins, namely α -actinin-1, α -actinin-2, α -actinin-3 and α -actinin-4, which are localized to muscle and non-muscle cells, including skeletal, cardiac and smooth muscle cells, as well as within the cytoskeleton. Each α -actinin protein contains one Actin-binding domain, two calponin-homology domains, two EF-hand domains and four spectrin repeats, through which they function as bundling proteins that can cross-link F-Actin, thus anchoring Actin to a variety of intracellular structures. Defects in the gene encoding α -actinin-4 are the cause of focal segmental glomerulosclerosis 1 (FSGS1), a common renal lesion characterized by decreasing kidney function and, ultimately, renal failure.

REFERENCES

1. Youssoufian, H., et al. 1990. Cloning and chromosomal localization of the human cytoskeletal α -actinin gene reveals linkage to the β -spectrin gene. *Am. J. Hum. Genet.* 47: 62-71.
2. Nishiyama, M., et al. 1990. Expression of human α -actinin in human hepatocellular carcinoma. *Cancer Res.* 50: 6291-6294.
3. Yürüker, B. and Niggli, V. 1992. α -actinin and vinculin in human neutrophils: reorganization during adhesion and relation to the Actin network. *J. Cell Sci.* 101: 403-414.
4. Knudsen, K.A., et al. 1995. Interaction of α -actinin with the cadherin/catenin cell-cell adhesion complex via α -catenin. *J. Cell Biol.* 130: 67-77.

CHROMOSOMAL LOCATION

Genetic locus: ACTN4 (human) mapping to 19q13.2.

PRODUCT

α -actinin-4 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see α -actinin-4 shRNA Plasmid (h): sc-43101-SH and α -actinin-4 shRNA (h) Lentiviral Particles: sc-43101-V as alternate gene silencing products.

For independent verification of α -actinin-4 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43101A, sc-43101B and sc-43101C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

α -actinin-4 siRNA (h) is recommended for the inhibition of α -actinin-4 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

α -actinin-4 (G-4): sc-390205 is recommended as a control antibody for monitoring of α -actinin-4 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor α -actinin-4 gene expression knockdown using RT-PCR Primer: α -actinin-4 (h)-PR: sc-43101-PR (20 μ l, 516 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Barbolina, M.V., et al. 2008. Motility-related actinin α -4 is associated with advanced and metastatic ovarian carcinoma. *Lab. Invest.* 88: 602-614.
2. Yoshii, H., et al. 2013. Increased expression of α -actinin-4 is associated with unfavorable pathological features and invasiveness of bladder cancer. *Oncol. Rep.* 30: 1073-1080.
3. Wang, M.C., et al. 2015. α -actinin 4 is associated with cancer cell motility and is a potential biomarker in non-small cell lung cancer. *J. Thorac. Oncol.* 10: 286-301.

RESEARCH USE

For research use only, not for use in diagnostic procedures.