

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Integrin α6 siRNA (h): sc-43129



The Power to Question

BACKGROUND

Integrins are heterodimers composed of noncovalently associated transmembrane α and β subunits. The 16 α and 8 β subunits heterodimerize to produce more than 20 different receptors. Most integrin receptors bind ligands that are components of the extracellular matrix, including Fibronectin, collagen and Vitronectin. Certain integrins can also bind to soluble ligands such as fibrinogen, or to counterreceptors on adjacent cells such as the intracellular adhesion molecules (ICAMs), leading to aggregation of cells. Ligands serve to cross-link or cluster integrins by binding to adjacent integrin receptors; both receptor clustering and ligand occupancy are necessary for the activation of integrin-mediated responses. In addition to mediating cell adhesion and cytoskeletal organization, integrins function as signaling receptors. Signals transduced by integrins play a role in many biological processes, including cell growth, differentiation, migration and apoptosis.

REFERENCES

- 1. Hynes, R.O. 1992. Integrins: versatility, modulation and signaling in cell adhesion. Cell 69: 11-25.
- 2. Miyamoto, S., et al. 1995. Synergistic roles for receptor occupancy and aggregation in integrin transmembrane function. Science 267: 883-885.
- 3. Clark, E.A. and Brugge, J.S. 1995. Integrins and signal transduction pathways: the road taken. Science 268: 233-239.

CHROMOSOMAL LOCATION

Genetic locus: ITGA6 (human) mapping to 2q31.1.

PRODUCT

Integrin $\alpha 6$ siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Integrin $\alpha 6$ shRNA Plasmid (h): sc-43129-SH and Integrin $\alpha 6$ shRNA (h) Lentiviral Particles: sc-43129-V as alternate gene silencing products.

For independent verification of Integrin $\alpha 6$ (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43129A, sc-43129B and sc-43129C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Integrin $\alpha 6$ siRNA (h) is recommended for the inhibition of Integrin $\alpha 6$ expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Integrin α 6 (F-6): sc-374057 is recommended as a control antibody for monitoring of Integrin α 6 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Integrin $\alpha6$ gene expression knockdown using RT-PCR Primer: Integrin $\alpha6$ (h)-PR: sc-43129-PR (20 μ I, 426 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

- 1. Smadja, D.M., et al. 2014. α 6-Integrin is required for the adhesion and vasculogenic potential of hemangioma stem cells. Stem Cells 32: 684-693.
- Moon, H.G., et al. 2014. CCN1 secretion and cleavage regulate the lung epithelial cell functions after cigarette smoke. Am. J. Physiol. Lung Cell. Mol. Physiol. 307: L326-L337.
- 3. Fisher, M.L., et al. 2016. Transglutaminase interaction with α 6/ β 4-Integrin stimulates YAP1-dependent Δ Np63 α stabilization and leads to enhanced cancer stem cell survival and tumor formation. Cancer Res. 76: 7265-7276.
- 4. Grun, D., et al. 2018. NRP-1 interacts with GIPC1 and α 6/ β 4-Integrins to increase YAP1/ Δ Np63 α -dependent epidermal cancer stem cell survival. Oncogene 37: 4711-4722.
- Golbert, D.C.F., et al. 2018. Small interference ITGA6 gene targeting in the human thymic epithelium differentially regulates the expression of immunological synapse-related genes. Cell Adh. Migr. 12: 152-167.
- 6. Gao, X., et al. 2018. Physiological stretch induced proliferation of human urothelial cells via Integrin α 6-FAK signaling pathway. Neurourol. Urodyn. 37: 2114-2120.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**