



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



Cytokeratin 14 siRNA (m): sc-43310

BACKGROUND

Cytokeratins comprise a diverse group of intermediate filament proteins (IFPs) that are expressed in pairs in both keratinized and non-keratinized epithelial tissue, where they constitute up to 85% of mature keratinocytes in the vertebrate epidermis. Cytokeratins play a critical role in differentiation and tissue specialization and function to maintain the overall structural integrity of epithelial cells. The α -helical coiled-coil dimers associate laterally end-to-end to form 10 nm diameter filaments. Cytokeratins are useful markers of tissue differentiation and, in addition, they aid in the characterization of malignant tumors. In Bowen's disease, the characteristic malignancy of the epidermis exhibits distinct expression patterns of Cytokeratin 14. The gene encoding human Cytokeratin 14 maps to chromosome 17q21.2. Mutations in this gene lead to epidermolysis bullosa simplex, an inherited skin disorder characterized by skin blistering due to basal keratinocyte fragility.

REFERENCES

- Rosenberg, M., et al. 1988. A group of type I keratin genes on human chromosome 17: characterization and expression. *Mol. Cell. Biol.* 8: 722-736.
- Bonifas, J.M., et al. 1991. Epidermolysis bullosa simplex: evidence in two families for keratin gene abnormalities. *Science* 254: 1202-1205.
- van der Velden, L.A., et al. 1993. Cytokeratin expression in normal and (pre)malignant head and neck epithelia: an overview. *Head Neck* 15: 133-146.
- Chan, Y., et al. 1994. A human keratin 14 "knockout": the absence of K14 leads to severe epidermolysis bullosa simplex and a function for an intermediate filament protein. *Genes Dev.* 8: 2574-2587.
- Marceau, N., et al. 1995. Cytokeratin expression, fibrillar organization and subtle function in liver cells. *Biochem. Cell Biol.* 73: 619-625.
- Fuchs, E. 1995. Keratins and the skin. *Annu. Rev. Cell Dev. Biol.* 11: 123-153.
- Chen, H., et al. 1995. Keratin 14 gene mutations in patients with epidermolysis bullosa simplex. *J. Invest. Dermatol.* 105: 629-632.

CHROMOSOMAL LOCATION

Genetic locus: Krt1-14 (mouse) mapping to 11 D.

PRODUCT

Cytokeratin 14 siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Cytokeratin 14 shRNA Plasmid (m): sc-43310-SH and Cytokeratin 14 shRNA (m) Lentiviral Particles: sc-43310-V as alternate gene silencing products.

For independent verification of Cytokeratin 14 (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43310A and sc-43310B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Cytokeratin 14 siRNA (m) is recommended for the inhibition of Cytokeratin 14 expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Cytokeratin 14 (LL001): sc-53253 is recommended as a control antibody for monitoring of Cytokeratin 14 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended:
 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Cytokeratin 14 gene expression knockdown using RT-PCR Primer: Cytokeratin 14 (m)-PR: sc-43310-PR (20 μ l). Annealing temperature for the primers should be 55- 60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.