



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

S-100 α chain siRNA (m): sc-43355

BACKGROUND

The family of EF-hand type Ca^{2+} -binding proteins includes calbindin (previously designated vitamin D-dependent Ca^{2+} -binding protein), S-100 α and β , calgranulins A (also designated MRP8), B (also designated MRP14) and C (S-100 like proteins), and the parvalbumin family members, including parvalbumin α and parvalbumin β (also designated oncomodulin). The S-100 protein is involved in the regulation of cellular processes such as cell cycle progression and differentiation. Research also indicates that the S-100 protein may function in the activation of Ca^{2+} induced Ca^{2+} release, inhibition of microtubule assembly and inhibition of protein kinase C mediated phosphorylation. Two S-100 subunits, sharing 60% sequence identity, have been described as S-100 α chain and S-100 β chain. Three S-100 dimeric forms have been characterized, differing in their subunit composition of either two α chains, two β chains or one α and one β chain. S-100 localizes to the cytoplasm and nuclei of astrocytes, Schwann's cells, ependymomas and astroglomas. S-100 is also detected in almost all benign naevi, malignant melanocytic tumours and in Langerhans cells in the skin. Calbindin, S-100 proteins and parvalbumin proteins are each expressed in neural tissues. In addition, S-100 α and β are present in a variety of other tissues and calbindin is present in intestine and kidney.

REFERENCES

- Pfytter, G.E., et al. 1987. Developmental and functional studies of parvalbumin and Calbindin D28K in hypothalamic neurons grown in serum-free medium. *J. Neurochem.* 49: 442-451.
- Heizmann, C.W. 1988. Calcium-binding proteins of the EF-type. *J. Cardiovasc. Pharmacol.* 5: S30-S37.
- Kagi, U., et al. 1988. Developmental appearance of the Ca^{2+} -binding proteins parvalbumin, Calbindin D28K, S-100 proteins and Calmodulin during testicular development in the rat. *Cell Tissue Res.* 252: 359-365.
- Zimmer, D.B., et al. 1991. Isolation of a rat S-100 α cDNA and distribution of its mRNA in rat tissues. *Brain Res. Bull.* 27: 157-162.
- Rickmann, M. and Wolff, J.R. 1995. S-100 protein expression in subpopulations of neurons of rat brain. *Neuroscience* 67: 977-991.
- Wang, Y.Z. and Christakos, S. 1995. Retinoic acid regulates the expression of the calcium binding protein, Calbindin D28K. *Mol. Endocrinol.* 9: 1510-1521.
- Muntener, M., et al. 1995. Increase of skeletal muscle relaxation speed by direct injection of parvalbumin cDNA. *Proc. Natl. Acad. Sci. USA* 92: 6504-6508.
- Hitomi, J., et al. 1996. A novel calcium-binding protein in amniotic fluid. CAAF1: its molecular cloning and tissue distribution. *J. Cell Sci.* 109: 805-815.
- Beaudeau, J., et al. 1999. Pathophysiologic aspects of S-100 β protein: a new biological marker of brain pathology. *Ann. Biol. Clin.* 57: 261-272.

CHROMOSOMAL LOCATION

Genetic locus: S100a1 (mouse) mapping to 3 F1.

PRODUCT

S-100 α chain siRNA (m) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see S-100 α chain shRNA Plasmid (m): sc-43355-SH and S-100 α chain shRNA (m) Lentiviral Particles: sc-43355-V as alternate gene silencing products.

For independent verification of S-100 α chain (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43355A and sc-43355B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

S-100 α chain siRNA (m) is recommended for the inhibition of S-100 α chain expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

S-100 α chain (4c4.9): sc-58837 is recommended as a control antibody for monitoring of S-100 α chain gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor S-100 α chain gene expression knockdown using RT-PCR Primer: S-100 α chain (m)-PR: sc-43355-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.