



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## KIF1C siRNA (h): sc-43372

### BACKGROUND

The kinesins constitute a large family of microtubule-dependent motor proteins, which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Individual kinesin members play crucial roles in cell division, intracellular transport, and membrane trafficking events including endocytosis and transcytosis. KIF1C is a member of the KIF1/Unc104 family of kinesin-like proteins, which are involved in the transport of mitochondria or synaptic vesicles in axons. Human KIF1C maps to chromosome 17p13.2 and encodes a predicted 1,103 amino acid protein with abundant expression in heart and skeletal muscle. Tyrosine phosphorylation is a putative regulator of KIF1C mediated retrograde transport of Golgi vesicles to the endoplasmic reticulum. KIF1C is capable of forming homodimers and can noncovalently associate with 14-3-3  $\beta$ ,  $\gamma$ ,  $\epsilon$  and  $\zeta$ . In mouse macrophages, KIF1C is required for anthrax lethal toxin resistance.

### REFERENCES

1. Hamm-Alvarez, S.F. 1998. Molecular motors and their role in membrane traffic. *Adv. Drug Deliv. Rev.* 29: 229-242.
2. Dörner, C., Ciossek, T., Müller, S., Moller, P.H., Ullrich, A. and Lammers, R. 1998. Characterization of KIF1C, a new kinesin-like protein involved in vesicle transport from the Golgi apparatus to the endoplasmic reticulum. *J. Biol. Chem.* 273: 20267-20275.
3. Cole, D.G. 1999. Kinesin-II, the heteromeric kinesin. *Cell. Mol. Life Sci.* 56: 217-226.
4. Dörner, C., Ullrich, A., Haring, H.U. and Lammers, R. 1999. The kinesin-like motor protein KIF1C occurs in intact cells as a dimer and associates with proteins of the 14-3-3 family. *J. Biol. Chem.* 274: 33654-33660.

### CHROMOSOMAL LOCATION

Genetic locus: KIF1C (human) mapping to 17p13.2.

### PRODUCT

KIF1C siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KIF1C shRNA Plasmid (h): sc-43372-SH and KIF1C shRNA (h) Lentiviral Particles: sc-43372-V as alternate gene silencing products.

For independent verification of KIF1C (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43372A, sc-43372B and sc-43372C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^{\circ}$  C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^{\circ}$  C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

KIF1C siRNA (h) is recommended for the inhibition of human expression in KIF1C cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KIF1C gene expression knockdown using RT-PCR Primer: KIF1C (h)-PR: sc-43372-PR (20  $\mu$ l). Annealing temperature for the primers should be  $55-60^{\circ}$  C and the extension temperature should be  $68-72^{\circ}$  C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.