



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Rap 2B siRNA (m): sc-43516

BACKGROUND

Ras oncogenes encode GTP-binding proteins that are capable of transforming immortalized cells in culture. Two Ras-related human genes, designated RAP1A and RAP1B, encode 95% homologous proteins that share a similar C-terminal Cys-Ali-Ali-Xaa sequence with Ras proteins and are ubiquitously expressed in mammalian tissues. The putative "effector" domain of Ras proteins, whose integrity is required for cell transformation as well as interaction with the putative effector protein GAP, is conserved in both Rap 1 proteins. It has been postulated that p21Rap 1 acts to interfere with Ras effector function by binding to Ras GAP. In fact it is known that p21Rap 1 binds to Ras GAP *in vitro* in a GTP dependent manner without affecting p21Rap 1 GTPase activity. The Rap 2 protein shares 60% identity with the Rap 1A protein and exhibits a carboxy terminal CAAX motif and two upstream cysteines similar to those of the H-Ras, K-Ras and N-Ras proteins. In contrast with Rap 1, overexpression of Rap 2 does not interfere with the Ras signaling pathway.

REFERENCES

1. Pizon, V., et al. 1988. Human cDNAs Rap 1 and Rap 2 homologous to the *Drosophila* gene Dras3 encode proteins closely related to Ras in the "effector" region. *Oncogene* 3: 201-204.
2. Pizon, V., et al. 1988. Nucleotide sequence of a human cDNA encoding a Ras-related protein (Rap 1B). *Nucl. Acids Res.* 16: 7719.
3. Culine, S., et al. 1989. Expression of the Ras-related Rap genes in human tumors. *Int. J. Cancer* 44: 990-994.
4. Kitayama, H., et al. 1989. A Ras-related gene with transformation suppressor activity. *Cell* 56: 77-84.
5. Kim, S., et al. 1990. Tissue and subcellular distributions of the Smg-21/Rap 1/Krev-1 proteins which are partly distinct from those of c-Ras p21s. *Mol. Cell. Biol.* 10: 2645-2652.

CHROMOSOMAL LOCATION

Genetic locus: Rap2b (mouse) mapping to 3 E1.

PRODUCT

Rap 2B siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Rap 2B shRNA Plasmid (m): sc-43516-SH and Rap 2B shRNA (m) Lentiviral Particles: sc-43516-V as alternate gene silencing products.

For independent verification of Rap 2B (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43516A, sc-43516B and sc-43516C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Rap 2B siRNA (m) is recommended for the inhibition of Rap 2B siRNA (m) expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Rap 2A/B/C (C-2): sc-515711 is recommended as a control antibody for monitoring of Rap 2B gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Rab 2B gene expression knockdown using RT-PCR Primer: Rap 2B (m)-PR: sc-43516-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.