

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



CCK-BR shRNA (h) Lentiviral Particles: sc-43671-V



The Power to Overtion

BACKGROUND

Gastrin is responsible for the stimulation of various digestive functions. In response to gastrin, the stomach mucosa produces and secretes hydrochloric acid, and the pancreas secretes digestive enzymes. Gastrin also stimulates smooth muscle contraction and increases blood circulation and water secretion in the stomach and intestine. In the brain, cholecystokinin (CCK) is a neurotransmitter that is involved in satiety, stress and anxiety. CCK is expressed in the gastrointestinal (GI) system as well as the central nervous system (CNS), which provides further evidence that CCK modulates food consumption. Both CCK and gastrin mediate their effects through two G protein coupled receptors, CCK-AR and CCK-BR. CCK preferentially binds CCK-AR with high affinity, whereas CCK-BR binds to gastrin and CCK with nearly equal affinities. The cholecystokinin receptors and their ligands are potential therapeutic targets for GI or CNS diseases.

REFERENCES

- 1. Koh, T.J., et al. 1995. Molecular cloning and sequencing of the murine gastrin gene. Biochem. Biophys. Res. Commun. 216: 34-41.
- 2. Yassin, R.R. 1999. Signaling pathways mediating gastrin's growth-promoting effects. Peptides 20: 885-898.
- Crespi, F., et al. 2000. Involvement of cholecystokinin within craving for cocaine: role of cholecystokinin receptor ligands. Expert Opin. Investig. Drugs 9: 2249-2258.
- 4. de Tullio, P., et al. 2000. Therapeutic and chemical developments of cholecystokinin receptor ligands. Expert Opin. Investig. Drugs 9: 129-146.
- Beglinger, C., et al. 2001. Loxiglumide, a CCK-A receptor antagonist, stimulates calorie intake and hunger feelings in humans. Am. J. Physiol. Regul. Integr. Comp. Physiol. 280: 1149-1154.

CHROMOSOMAL LOCATION

Genetic locus: CCKBR (human) mapping to 11p15.4.

PRODUCT

CCK-BR shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0 x 10⁶ infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see CCK-BR siRNA (h): sc-43671 and CCK-BR shRNA Plasmid (h): sc-43671-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80 $^{\circ}$ C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4 $^{\circ}$ C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

CCK-BR shRNA (h) Lentiviral Particles is recommended for the inhibition of CCK-BR expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0 x 10 6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

CCK-BR (E-3): sc-166690 is recommended as a control antibody for monitoring of CCK-BR gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CCK-BR gene expression knockdown using RT-PCR Primer: CCK-BR (h)-PR: sc-43671-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**