



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GILZ shRNA (h) Lentiviral Particles: sc-43805-V

BACKGROUND

Glucocorticoid-induced leucine zipper (GILZ) is a leucine zipper protein expressed in normal lymphocytes from thymus, spleen and lymph nodes. It is absent in nonlymphoid tissues including brain, liver and kidney. GILZ mediates the immunosuppressive effects of glucocorticoid hormones; its expression is induced in T cells by dexamethasone. GILZ protects T cells from an anti-CD3 antibody-induced apoptosis by inhibiting Fas and Fas ligand expression. It interferes with Egr-2, Egr-3, NFAT/AP-1-inducible transcription factors and AP-1. The interaction of GILZ with c-Fos and c-Jun inhibits the binding of active AP-1 to its DNA consensus site *in vitro*. GILZ also binds NFκB subunits and inhibits the NFκB nuclear translocation. It inhibits T cell receptor-induced interleukin-2/interleukin-2 receptor expression. The binding of GILZ to Raf-1 prevents Raf-MEK-ERK activation in the MAPK pathway. GILZ is expressed by normal macrophages in nonlymphoid tissues and by tumor-infiltrating macrophages in Burkitt lymphomas. The gene encoding human GILZ maps to chromosome Xq22.3.

REFERENCES

1. D'Adamo, F., et al. 1997. A new dexamethasone-induced gene of the leucine zipper family protects T lymphocytes from TCR/CD3-activated cell death. *Immunity* 7: 803-812.
2. Mittelstadt, P.R., et al. 2001. Inhibition of AP-1 by the glucocorticoid-inducible protein GILZ. *J. Biol. Chem.* 276: 29603-29610.
3. Ayroldi, E., et al. 2001. Modulation of T-cell activation by the glucocorticoid-induced leucine zipper factor via inhibition of nuclear factor κB. *Blood* 98: 743-753.
4. Cannarile, L., et al. 2001. Cloning, chromosomal assignment and tissue distribution of human GILZ, a glucocorticoid hormone-induced gene. *Cell Death Differ.* 8: 201-203.

CHROMOSOMAL LOCATION

Genetic locus: TSC22D3 (human) mapping to Xq22.3.

PRODUCT

GILZ shRNA (h) Lentiviral Particles are concentrated, transduction-ready viral particles containing a target-specific construct that encodes a 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μl frozen stock containing 1.0 x 10⁶ infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see GILZ siRNA (h): sc-43805 and GILZ shRNA Plasmid (h): sc-43805-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

GILZ shRNA (h) Lentiviral Particles is recommended for the inhibition of GILZ expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μl frozen viral stock containing 1.0 x 10⁶ infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

GILZ (N-18): sc-26518 is recommended as a control antibody for monitoring of GILZ gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker™ compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GILZ gene expression knockdown using RT-PCR Primer: GILZ (h)-PR: sc-43805-PR (20 μl, 414 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.