



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

group II sPLA₂ siRNA (m): sc-43818

BACKGROUND

Phospholipases catalyze the release of fatty acids from phospho-lipids. One member of the phospholipase family, iPLA₂, is detected as a membrane-bound protein with multiple smaller isoforms, which result from alternative splicing. Two isoforms, ankyrin- iPLA₂-1 and 2, lack the catalytic domain and are thought to be involved in the negative regulation of iPLA₂ activity. The SH-iPLA₂ isoform is cytoplasmic-ally localized. The human gene encoding iPLA₂ maps to chromosome 22q13.1. Another phospholipase, sPLA₂, belongs to a family of secretory phospholipases A₂, which represent an expanding family of related enzymes. sPLA₂ has both membrane bound and secreted forms that are encoded by a single gene. sPLA₂ is involved in the regulation of phospholipid metabolism in biomembranes and in eicosanoid biosynthesis.

REFERENCES

1. Scott, D.L., et al. 1991. Structures of free and inhibited human secretory phospholipase A₂ from inflammatory exudate. *Science* 254: 1007-1010.
2. Lehninger, A., et al. 1993. Principles of Biochemistry Second Edition. New York: Worth Publishers.
3. Cupillard, L., et al. 1997. Cloning, chromosomal mapping, and expression of a novel human secretory phospholipase A₂. *J. Biol. Chem.* 272: 15745-15752.
4. Kitadokoro, K., et al. 1998. Crystal structure of human secretory phospholipase A₂-IIA complex with the potent indolizine inhibitor 120-1032. *J. Biochem.* 123: 619-623.
5. Ma, Z., et al. 1999. Human pancreatic islets express mRNA species encoding two distinct catalytically active isoforms of group VI phospholipase A₂ (iPLA₂) that arise from an exon-skipping mechanism of alternative splicing of the transcript from the iPLA₂ gene on chromosome 22q13.1. *J. Biol. Chem.* 274: 9607-9616.
6. Larsson-Forsell, P.K., et al. 1999. The human calcium-independent phospholipase A₂ gene multiple enzymes with distinct properties from a single gene. *Eur. J. Biochem.* 262: 575-585.

CHROMOSOMAL LOCATION

Genetic locus: Pla2g2a (mouse) mapping to 4 D3.

PRODUCT

group II sPLA₂ siRNA (m) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see group II sPLA₂ shRNA Plasmid (m): sc-43818-SH and group II sPLA₂ shRNA (m) Lentiviral Particles: sc-43818-V as alternate gene silencing products.

For independent verification of group II sPLA₂ (m) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43818A, sc-43818B and sc-43818C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

group II sPLA₂ siRNA (m) is recommended for the inhibition of group II sPLA₂ expression in mouse cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor group II sPLA₂ gene expression knockdown using RT-PCR Primer: group II sPLA₂ (m)-PR: sc-43818-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.