

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



group VI iPLA₂ shRNA (h) Lentiviral Particles: sc-43819-V



The Power to Question

BACKGROUND

Phospholipases catalyze the release of fatty acids from phospholipids. One member of the phospholipase family, iPLA2, is detected as a membrane-bound protein with multiple smaller isoforms, which result from alternative splicing. Two isoforms, Ankyrin-iPLA2-1 and -2, lack the catalytic domain and are thought to be involved in the negative regulation of iPLA2 activity. The SH-iPLA2 isoform is cytoplasmically localized. The human gene encoding iPLA2 maps to chromosome 22q13.1. Another phopholipase, sPLA2, belongs to a family of secretory phospholipases A2, which represent an expanding family of related enzymes. sPLA2 has both membrane bound and secreted forms that are encoded by a single gene. sPLA2 is involved in the regulation of phospholipid metabolism in biomembranes and in eicosanoid biosynthesis.

REFERENCES

- 1. Scott, D.L., et al. 1991. Structures of free and inhibited human secretory phospholipase A₂ from inflammatory exudate. Science 254: 1007-1010.
- Lehninger, A., et al. 1993. Principles of biochemistry second edition. Worth Publishers.
- Cupillard, L., et al. 1997. Cloning, chromosomal mapping, and expression of a novel human secretory phospholipase A₂. J. Biol. Chem. 272: 15745-15752.
- 4. Kitadokoro, K., et al. 1998. Crystal structure of human secretory phospholipase A_2 -IIA complex with the potent indolizine inhibitor 120-1032. J. Biochem. 123: 619-623.
- 5. Ma, Z., et al. 1999. Human pancreatic islets express mRNA species encoding two distinct catalytically active isoforms of group VI phospholipase A₂ (iPLA₂) that arise from an exon-skipping mechanism of alternative splicing of the transcript from the iPLA₂ gene on chromosome 22q13.1. J. Biol. Chem. 274: 9607-9616.
- Larsson-Forsell, P.K., et al. 1999. The human calcium-independent phospholipase A₂ gene multiple enzymes with distinct properties from a single gene. Eur. J. Biochem. 262: 575-585.

CHROMOSOMAL LOCATION

Genetic locus: PLA2G6 (human) mapping to 22q13.1.

PRODUCT

group VI iPLA $_2$ shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0 x 10⁶ infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see group VI iPLA $_2$ siRNA (h): sc-43819 and group VI iPLA $_2$ shRNA Plasmid (h): sc-43819-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

group VI iPLA $_2$ shRNA (h) Lentiviral Particles is recommended for the inhibition of group VI iPLA $_2$ expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0 x 10 6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

group VI iPLA $_2$ (D-4): sc-376563 is recommended as a control antibody for monitoring of group VI iPLA $_2$ gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluo-rescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor group VI iPLA $_2$ gene expression knockdown using RT-PCR Primer: group VI iPLA $_2$ (h)-PR: sc-43819-PR (20 μ I, 428 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.