



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

IEX-1 siRNA (h): sc-43859

BACKGROUND

Tumors are frequently observed as resistant to apoptotic induction by FAS, tumor necrosis factor α (TNF α) or irradiation. This anti-death activity may be attributed to immediate early-response genes that are regulated at the transcriptional level, including the protein IEX-1. IEX-1 was originally characterized as a gene induced by ultraviolet radiation and TNF α , which protected human squamous carcinoma cells from apoptosis. Subsequent transfection studies have also shown that expression of IEX-1 in human keratinocytes and mouse fibroblasts results in more rapid proliferation of the cells as compared with controls. The promoter region of IEX-1 contains binding motifs for both NF κ B and p53, suggesting that these proteins may regulate its expression. IEX-1 is expressed as both a longer form (IEX-1_L) and a splice variant, designated IEX-1_S to indicate the shorter form and it is localized to the nucleus and perinuclear region.

REFERENCES

1. Beyaert, R., et al. 1994. Molecular mechanisms of tumor necrosis factor-induced cytotoxicity. What we do understand and what we do not. *FEBS Lett.* 340: 9-16.
2. Kondratyev, A.D., et al. 1996. Identification and characterization of a radiation-inducible glycosylated human early-response gene. *Cancer Res.* 56: 1498-1502.
3. Wu, M.X., et al. 1998. IEX-1L, an apoptosis inhibitor involved in NF κ B-mediated cell survival. *Science* 281: 998-1001.
4. Kobayashi, T., et al. 1998. Regulation of a novel immediate early response gene, IEX-1, in keratinocytes by 1,25-dihydroxyvitamin D₃. *Biochem. Biophys. Res. Commun.* 251: 868-873.
5. Kumar, R., et al. 1998. A novel immediate early response gene, IEX-1, is induced by ultraviolet radiation in human keratinocytes. *Biochem. Biophys. Res. Commun.* 253: 336-341.
6. Schafer, H., et al. 1998. The promoter of human p22/PACAP response gene 1 (PRG1) contains functional binding sites for the p53 tumor suppressor and for NF κ B. *FEBS Lett.* 436: 139-143.

CHROMOSOMAL LOCATION

Genetic locus: IER3 (human) mapping to 6p21.33.

PRODUCT

IEX-1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see IEX-1 shRNA Plasmid (h): sc-43859-SH and IEX-1 shRNA (h) Lentiviral Particles: sc-43859-V as alternate gene silencing products.

For independent verification of IEX-1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-43859A, sc-43859B and sc-43859C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

IEX-1 siRNA (h) is recommended for the inhibition of IEX-1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor IEX-1 gene expression knockdown using RT-PCR Primer: IEX-1 (h)-PR: sc-43859-PR (20 μ l, 500 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Emma, M.R., et al. 2016. NUPR1, a new target in liver cancer: implication in controlling cell growth, migration, invasion and sorafenib resistance. *Cell Death Dis.* 7: e2269.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.