

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

LTA4H shRNA (h) Lentiviral Particles: sc-43895-V



BACKGROUND

Leukotrienes are biologically active compounds formed from arachidonic acid or polyunsaturated fatty acids that are important in host defense reactions and play a pathophysiological role in inflammation and allergic reactions. LTA4H (leukotriene A4-hydrolase) is a Zn²⁺-containing enzyme with both epoxide hydrolase and aminopeptidase activity. As an epoxide hydrolase, LTA4H catalyzes the hydration of LTA4 to leukotriene B4 (LTB4, 5S,12R-dihydroxy-6,14-cis-8,10-trans-eicosatetraenoic acid), a potent lipid chemoattractant that influences inflammation, immune responses and host defense against infection. As an aminopeptidase, LTA4H catalyzes the cleavage of amides of paranitroaniline. The human LTA4H gene encodes a 610 amino acid protein.

REFERENCES

- 1. Minami, M., Ohno, S., Kawasaki, H., Radmark, O., Samuelsson, B., Jornvall, H., Shimizu, T., Seyama, Y. and Suzuki, K. 1987. Molecular cloning of a cDNA coding for human leukotriene A4 hydrolase. Complete primary structure of an enzyme involved in eicosanoid synthesis. J. Biol. Chem. 262: 13873-13876.
- 2. Gierse, J.K., Luckow, V.A., Askonas, L.J., Duffin, K.L., Aykent, S., Bild, G.S., Rodi, C.P., Sullivan, P.M., Bourner, M.J., Kimack, N.M., et al. 1993. Highlevel expression and purification of human leukotriene A4 hydrolase from insect cells infected with a baculovirus vector. Protein Expr. Purif. 4: 358-366.
- 3. Parnas, B.L., Durley, R.C., Rhoden, E.E., Kilpatrick, B.F., Makkar, N., Thomas, K.E., Smith, W.G. and Corley, D.G. 1996. Isolation and structure of leukotriene-A4 hydrolase inhibitor: 8(S)-amino-2(R)-methyl-7-oxononanoic acid produced by Streptomyces diastaticus. J. Nat. Prod. 59: 962-964.
- 4. Online Mendelian Inheritance in Man, OMIM™. 1998. Johns Hopkins University, Baltimore, MD. MIM Number: 151570. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- 5. Thunnissen, M.M., Nordlund, P. and Haeggstrom, J.Z. 2001. Crystal structure of human leukotriene A4 hydrolase, a bifunctional enzyme in inflammation. Nat. Struct. Biol. 8: 131-135.

CHROMOSOMAL LOCATION

Genetic locus: LTA4H (human) mapping to 12q23.1.

PRODUCT

LTA4H shRNA (h) Lentiviral Particles is a pool of concentrated, transductionready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 µl frozen stock containing 1.0 x 10⁶ infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see LTA4H siRNA (h): sc-43895 and LTA4H shRNA Plasmid (h): sc-43895-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

LTA4H shRNA (h) Lentiviral Particles is recommended for the inhibition of LTA4H expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 µl frozen viral stock containing 1.0 x 10⁶ infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

LTA4H (D-6): sc-390567 is recommended as a control antibody for monitoring of LTA4H gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker[™] compatible goat antimouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LTA4H gene expression knockdown using RT-PCR Primer: LTA4H (h)-PR: sc-43895-PR (20 µl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.