

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

SANTA CRUZ BIOTECHNOLOGY, INC.

Msx-1 shRNA (h) Lentiviral Particles: sc-43945-V



BACKGROUND

Msx homeobox genes encode for transcription factors that control morphogenesis and are expressed at sites of epithelial-mesenchymal interaction during embryogenesis, such as the tooth. Two of these genes, Msx-1 and Msx-2, are key factors for the development of tooth and craniofacial skeleton. Msx-1 also downregulates a master gene of skeletal cells differentiation. Msx-1 and Msx-2 contribute to the initial patterning of dentition as well as playing a pivotal role in terminal cell differentiation. In addition, Msx-1 and Msx-2 are expressed in the epidermis, hair follicles and fibroblasts of the developing fetal skin. In adult skin, Msx-1 and Msx-2 expression is confined to epithelially derived structures. Msx-2 is detected as a diffuse cytoplasmic signal in fetal epidermis and portions of the hair follicle and dermis, but is localized to the nucleus in the adult epidermis. Msx-1 and Msx-2 are also expressed during critical developmental stages of neural tube and neural crest, suggesting that these genes play an important role in organogenesis.

REFERENCES

- 1. Lezot, F., et al. 2000. Biomineralization, life-time of odontogenic cells and differential expression of the two homeobox genes Msx1 and Dxl2 in transgenic mice. J. Bone Miner. Res. 15: 430-441.
- 2. Maas, R. and Bei, M. 1997. The genetic control of early tooth development. Crit. Rev. Oral Biol. 8: 4-39.
- 3. Blin-Wakkach, C., et al. 2001. Endogenous Msx-1 antisense transcript: In vivo and in vitro evidences, structure, and potential involvement in skeleton development in mammals. Proc. Natl. Acad. Sci. USA 98: 7336-7341.
- 4. Stelnicki, E.J., et al. 1997. The human homeobox genes MSX1, MSX2, and MOX1 are differentially expressed in the dermis and epidermis in fetal and adult skin. Differentiation 62: 33-41.
- 5. Foerst-Potts, L. and Sadler, T.W. 1997. Disruption of Msx-1 and Msx-2 reveals roles for these genes in craniofacial, eye, and axial development. Dev. Dyn. 209: 70-84.

CHROMOSOMAL LOCATION

Genetic locus: MSX1 (human) mapping to 4p16.2.

PRODUCT

Msx-1 shRNA (h) Lentiviral Particles is a pool of concentrated, transductionready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 µl frozen stock containing 1.0 x 10⁶ infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Msx-1 siRNA (h): sc-43945 and Msx-1 shRNA Plasmid (h): sc-43945-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

APPLICATIONS

Msx-1 shRNA (h) Lentiviral Particles is recommended for the inhibition of Msx-1 expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 µl frozen viral stock containing 1.0 x 10⁶ infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Msx-1 (E-20): sc-17727 is recommended as a control antibody for monitoring of Msx-1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz Marker[™] compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruz[™] Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Msx-1 gene expression knockdown using RT-PCR Primer: Msx-1 (h)-PR: sc-43945-PR (20 µl, 527 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.