



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PLC ϵ siRNA (h): sc-44024

BACKGROUND

Phosphoinositide-specific phospholipase C (PLC) plays a crucial role in the initiation of receptor mediated signal transduction through the generation of the two second messengers, inositol 1,4,5-triphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. There are many mammalian PLC isozymes, including PLC β 1, PLC β 2, PLC β 3, PLC β 4, PLC γ 1, PLC γ 2, PLC δ 1, PLC δ 2 and PLC ϵ . Phospholipase C ϵ (PLC ϵ) is characterized by possession of CDC25 homology and Ras/Rap1-associating domains. PLC ϵ is translocated from the cytoplasm to the plasma membrane and activated by direct association with Ras at its Ras/Rap1-associating domain.

REFERENCES

1. Rhee, S.G. and Choi, K.D. 1992. Regulation of inositol phospholipid-specific phospholipase C isozymes. *J. Biol. Chem.* 267: 12393-12396.
2. Kelley, G.G., et al. 2001. Phospholipase C ϵ : a novel Ras effector. *EMBO J.* 20: 743-754.
3. Jin, T.G., et al. 2001. Role of the CDC25 homology domain of phospholipase C ϵ in amplification of Rap1-dependent signaling. *J. Biol. Chem.* 276: 30301-30307.
4. Wing, M.R., et al. 2001. Activation of phospholipase C- ϵ by heterotrimeric G protein $\beta\gamma$ -subunits. *J. Biol. Chem.* 276: 48257-48261.
5. Song, C., et al. 2002. Differential roles of Ras and Rap1 in growth factor-dependent activation of phospholipase C ϵ . *Oncogene* 21: 8105-8113.
6. Wu, D., et al. 2003. Neuronal lineage-specific induction of phospholipase C ϵ expression in the developing mouse brain. *Eur. J. Neurosci.* 17: 1571-1580.
7. Wing, M.R., et al. 2003. Direct activation of phospholipase C- ϵ by Rho. *J. Biol. Chem.* 278: 41253-41258.
8. Wing, M.R., et al. 2003. PLC ϵ : a shared effector protein in Ras-, Rho-, and G α β γ -mediated signaling. *Mol. Interv.* 3: 273-280.
9. Seifert, J.P., et al. 2004. RhoA activates purified phospholipase C- ϵ by a guanine nucleotide-dependent mechanism. *J. Biol. Chem.* 279: 47992-47997.

CHROMOSOMAL LOCATION

Genetic locus: PLCE1 (human) mapping to 10q23.33.

PRODUCT

PLC ϵ siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PLC ϵ shRNA Plasmid (h): sc-44024-SH and PLC ϵ shRNA (h) Lentiviral Particles: sc-44024-V as alternate gene silencing products.

For independent verification of PLC ϵ (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44024A, sc-44024B and sc-44024C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PLC ϵ siRNA (h) is recommended for the inhibition of PLC ϵ expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PLC ϵ gene expression knockdown using RT-PCR Primer: PLC ϵ (h)-PR: sc-44024-PR (20 μ l, 427 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

SELECT PRODUCT CITATIONS

1. Hashimoto, A., et al. 2015. Cilostazol induces PGI2 production via activation of the downstream Epac-1/Rap 1 signaling cascade to increase intracellular calcium by PLC ϵ and to activate p44/42 MAPK in human aortic endothelial cells. *PLoS ONE* 10: e0132835.
2. Li, Y. and Luan, C. 2018. PLCE1 promotes the invasion and migration of esophageal cancer cells by up-regulating the PKC α /NF κ B pathway. *Yonsei Med. J.* 59: 1159-1165.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.