



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## PTP $\gamma$ siRNA (h): sc-44049

### BACKGROUND

Protein-tyrosine phosphatases, or PTPs, are type I transmembrane proteins, membrane associated proteins or proteins localized in nuclei. Examples of transmembrane PTPs are LAR, PTP $\alpha$ , PTP $\beta$ , PTP $\gamma$ , PTP $\delta$ , PTP $\epsilon$ , PTP $\zeta$ , PTP $\kappa$  and PTP $\mu$ . Transmembrane PTPs play diverse roles during development and in adult tissues. Immunodepletion studies have suggested LAR to be a regulator of Insulin receptor phosphorylation. PTP $\alpha$  activity is increased twofold in response to phorbol ester stimulation, resulting in serine phosphorylation either directly or indirectly by members of the PKC family. Overexpression of v-H-Ras and Neu, but not Myc or Int2, in mammary tumors has been shown to induce PTP $\epsilon$  expression. An alternative splicing event leads to a nervous tissue-specific chondroitin sulfate proteoglycan called phosphacan, which represents the amino-terminal portion of PTP $\zeta$ . PTP $\kappa$  and PTP $\mu$  share a conserved amino terminal 160 amino acid MAM domain which facilitates homophilic binding. PTP $\mu$  localizes to points of cell contact and may be involved in regulating the assembly and disassembly of cadherin/catenin complexes *in vivo*.

### REFERENCES

- Ahmad, F., et al. 1995. Increased abundance of the receptor-type protein-tyrosine phosphatase LAR accounts for the elevated Insulin receptor dephosphorylating activity in adipose tissue of obese human subjects. *J. Clin. Invest.* 95: 2806-2812.
- den Hertog, J., et al. 1995. Stimulation of receptor protein-tyrosine phosphatase  $\alpha$  activity and phosphorylation by phorbol ester. *Cell Growth Differ.* 6: 303-307.
- Elson, A., et al. 1995. Protein-tyrosine phosphatase  $\epsilon$ . An isoform specifically expressed in mouse mammary tumors initiated by v-H-Ras or Neu. *J. Biol. Chem.* 270: 26116-26122.
- Brady-Kalnay, S.M., et al. 1995. Receptor protein-tyrosine phosphatase PTP $\mu$  associates with cadherins and catenins *in vivo*. *J. Cell Biol.* 130: 977-986.
- Zondag, G.C., et al. 1995. Homophilic interactions mediated by receptor tyrosine phosphatases  $\mu$  and  $\kappa$ . A critical role for the novel extracellular MAM domain. *J. Biol. Chem.* 270: 14247-14250.

### CHROMOSOMAL LOCATION

Genetic locus: PTPRG (human) mapping to 3p14.2.

### PRODUCT

PTP $\gamma$  siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PTP $\gamma$  shRNA Plasmid (h): sc-44049-SH and PTP $\gamma$  shRNA (h) Lentiviral Particles: sc-44049-V as alternate gene silencing products.

For independent verification of PTP $\gamma$  (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44049A, sc-44049B and sc-44049C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

PTP $\gamma$  siRNA (h) is recommended for the inhibition of PTP $\gamma$  expression in human cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PTP $\gamma$  gene expression knockdown using RT-PCR Primer: PTP $\gamma$  (h)-PR: sc-44049-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.