



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Serglycin shRNA (h) Lentiviral Particles: sc-44093-V

BACKGROUND

Proteoglycans stored in the secretory granules of many hematopoietic cells contain a protease-resistant peptide core that may be important for neutralizing hydrolytic enzymes. Serglycin is associated with the macromolecular complex of granzymes and perforin, which may serve as a mediator of granule-mediated apoptosis. Serglycin is a chondroitin sulfate-bearing proteoglycan that functions in the transport of cationic granular proteins. The immune system relies on granule exocytosis as the main pathway for elimination of virus-infected cells and tumor cells by cytotoxic T lymphocytes and natural killer cells, thus indicating an important role for Serglycin in normal immune function.

REFERENCES

1. Raja, S.M., et al. 2002. Cytotoxic cell granule-mediated apoptosis. Characterization of the macromolecular complex of granzyme B with Serglycin. *J. Biol. Chem.* 277: 49523-49530.
2. Lemansky, P., et al. 2003. Targeting myeloperoxidase to azurophilic granules in HL-60 cells. *J. Leukoc. Biol.* 74: 542-550.
3. Lieberman, J. 2003. The ABCs of granule-mediated cytotoxicity: new weapons in the arsenal. *Nat. Rev. Immunol.* 3: 361-370.
4. Schick, B.P., et al. 2003. Serglycin proteoglycan expression and synthesis in embryonic stem cells. *Biochim. Biophys. Acta* 1593: 259-267.
5. Abrink, M., et al. 2004. Serglycin is essential for maturation of mast cell secretory granule. *J. Biol. Chem.* 279: 40897-40905.
6. Niemann, C.U., et al. 2004. Localization of Serglycin in human neutrophil granulocytes and their precursors. *J. Leukoc. Biol.* 76: 406-415.

CHROMOSOMAL LOCATION

Genetic locus: SRGN (human) mapping to 10q22.1.

PRODUCT

Serglycin shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see Serglycin siRNA (h): sc-44093 and Serglycin shRNA Plasmid (h): sc-44093-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80°C . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4°C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

Serglycin shRNA (h) Lentiviral Particles is recommended for the inhibition of Serglycin expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

Serglycin (C-11): sc-374657 is recommended as a control antibody for monitoring of Serglycin gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Serglycin gene expression knockdown using RT-PCR Primer: Serglycin (h)-PR: sc-44093-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ\text{C}$ and the extension temperature should be $68-72^\circ\text{C}$.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.