



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

SRp55 shRNA (h) Lentiviral Particles: sc-44118-V

BACKGROUND

Pre-mRNA splicing is a critical step in the post-transcriptional regulation of gene expression. Several protein complexes are involved in proper mRNA splicing and transport. Serine/arginine-rich (SR) proteins SRp55, SRp30c, and htra2 β 1 regulate exon 2 and 10 splicing. The first two inhibit both exons and SRp55 also plays a role in exon inclusion after the removal of intronic splicing silencer sequences. SRp55 plays a major role in maintaining normal FGFR1 α -exon inclusion.

REFERENCES

1. Ring, H.Z. and Lis, J.T. 1994. The SR protein B52/SRp55 is essential for *Drosophila* development. *Mol. Cell. Biol.* 14: 7499-7506.
2. Nagel, R.J., et al. 1998. Specific binding of an exonic splicing enhancer by the pre-mRNA splicing factor SRp55. *RNA* 4: 11-23.
3. Lemaire, R., et al. 1999. SF2 and SRp55 regulation of CD45 exon 4 skipping during T cell activation. *Eur. J. Immunol.* 29: 823-837.
4. Tran, Q., et al. 2003. Human transformer 2 β and SRp55 interact with a calcitonin-specific splice enhancer. *Biochim. Biophys. Acta* 1625: 141-152.
5. Tran, Q., et al. 2003. SRp55 is a regulator of calcitonin/CGRP alternative RNA splicing. *Biochemistry* 42: 951-957.
6. Lai, M.C., et al. 2003. Differential effects of hyperphosphorylation on splicing factor SRp55. *Biochem. J.* 371 (Pt. 3): 937-945.
7. Jin, W. et al. 2004. Enhancer-dependent splicing of FGFR1 α -exon is repressed by RNA interference-mediated down-regulation of SRp55. *Cancer. Res.* 64: 8901-8905.

CHROMOSOMAL LOCATION

Genetic locus: SFRS6 (human) mapping to 20q13.11.

PRODUCT

SRp55 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see SRp55 siRNA (h): sc-44118 and SRp55 shRNA Plasmid (h): sc-44118-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80 $^{\circ}$ C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4 $^{\circ}$ C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

SRp55 shRNA (h) Lentiviral Particles is recommended for the inhibition of SRp55 expression in human cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

SRp55 (G-15): sc-34198 is recommended as a control antibody for monitoring of SRp55 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use donkey anti-goat IgG-HRP: sc-2020 (dilution range: 1:2000-1:100,000) or Cruz MarkerTM compatible donkey anti-goat IgG-HRP: sc-2033 (dilution range: 1:2000-1:5000), Cruz MarkerTM Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use donkey anti-goat IgG-FITC: sc-2024 (dilution range: 1:100-1:400) or donkey anti-goat IgG-TR: sc-2783 (dilution range: 1:100-1:400) with UltraCruzTM Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SRp55 gene expression knockdown using RT-PCR Primer: SRp55 (h)-PR: sc-44118-PR (20 μ l, 597 bp). Annealing temperature for the primers should be 55-60 $^{\circ}$ C and the extension temperature should be 68-72 $^{\circ}$ C.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.