



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## TRPM5 siRNA (h): sc-44169

### BACKGROUND

Transient receptor potential ion channels (TRPCs) are a superfamily of six transmembrane segment-spanning, gated cation channels. TRPC subtypes mediate store-operated  $\text{Ca}^{2+}$  entry, a process involving  $\text{Ca}^{2+}$  influx and replenishment of  $\text{Ca}^{2+}$  stores formerly emptied through the action of inositol 1,4,5-trisphosphate production and other  $\text{Ca}^{2+}$  mobilizing agents. TRP ion channels influence calcium-depletion induced calcium influx processes in response to chemo-, mechano- and osmoregulatory events. TRPM5 forms a cation channel that is directly activated by micromolar concentrations of intracellular  $\text{Ca}^{2+}$ . Sustained exposure to  $\text{Ca}^{2+}$  desensitizes TRPM5 channels, but phosphatidylinositol-4,5-bisphosphate reverses desensitization, partially restoring channel activity. TRPM5 channels are nonselective among monovalent cations and not detectably permeable to divalent cations.

### REFERENCES

- Liu, D. and Liman, E.R. 2003. Intracellular  $\text{Ca}^{2+}$  and the phospholipid PIP2 regulate the taste transduction ion channel TRPM5. *Proc. Natl. Acad. Sci. USA* 100: 15160-15165.
- Prawitt, D., Monteilh-Zoller, M.K., Brixel, L., Spangenberg, C., Zabel, B., Fleig, A. and Penner, R. 2003. TRPM5 is a transient  $\text{Ca}^{2+}$ -activated cation channel responding to rapid changes in  $[\text{Ca}^{2+}]_i$ . *Proc. Natl. Acad. Sci. USA* 100: 15166-15171.
- Hofmann, T., Chubanov, V., Gudermann, T. and Montell, C. 2003. TRPM5 is a voltage-modulated and  $\text{Ca}^{2+}$ -activated monovalent selective cation channel. *Curr. Biol.* 13: 1153-1158.
- Perez, C.A., Margolskee, R.F., Kinnamon, S.C. and Ogura, T. 2003. Making sense with TRP channels: store-operated calcium entry and the ion channel TRPM5 in taste receptor cells. *Cell Calcium* 33: 541-549.
- Perraud A.L., Knowles, H.M. and Schmitz, C. 2004. Novel aspects of signaling and ion-homeostasis regulation in immunocytes. The TRPM ion channels and their potential role in modulating the immune response. *Mol. Immunol.* 41: 657-673.

### CHROMOSOMAL LOCATION

Genetic locus: TRPM5 (human) mapping to 11p15.5.

### PRODUCT

TRPM5 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu\text{M}$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TRPM5 shRNA Plasmid (h): sc-44169-SH and TRPM5 shRNA (h) Lentiviral Particles: sc-44169-V as alternate gene silencing products.

For independent verification of TRPM5 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44169A, sc-44169B and sc-44169C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at  $-20^\circ\text{C}$  with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at  $-20^\circ\text{C}$ , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu\text{l}$  of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu\text{l}$  of RNase-free water makes a 10  $\mu\text{M}$  solution in a 10  $\mu\text{M}$  Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

TRPM5 siRNA (h) is recommended for the inhibition of TRPM5 expression in human cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu\text{M}$  in 66  $\mu\text{l}$ . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TRPM5 gene expression knockdown using RT-PCR Primer: TRPM5 (h)-PR: sc-44169-PR (20  $\mu\text{l}$ ). Annealing temperature for the primers should be  $55-60^\circ\text{C}$  and the extension temperature should be  $68-72^\circ\text{C}$ .

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.