



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

ACP1 shRNA (m) Lentiviral Particles: sc-44359-V

BACKGROUND

Regulation of intracellular concentrations of flavoenzymes and flavin coenzymes is essential for proper cell homeostasis. Red cell acid phosphatase, known as ACP1, catalyzes the transfer of phosphate from phosphate ester substrates to suitable acceptor alcohols such as methanol and glycerol. ACP1 is a genetically polymorphic, cytoplasmic low-molecular-weight flavin mononucleotide phosphatase that regulates the intracellular concentrations of flavin coenzymes. The human ACP1 gene maps to chromosome 2p25 and encodes a pair of isozymes, Bf (α) and Bs (β). The ACP1 α and β isozymes are not glycosylated. Both ACP1- α and ACP1- β isozymes are 157 amino acids in length; however the two forms differ in sequence over an internal 34 residue segment. The two isoforms are believed to differ in substrate specificity.

REFERENCES

1. Golden, V.L., et al. 1986. Phenotypic variation in the phosphotransferase activity of human red cell acid phosphatase (ACP1). *Hum. Genet.* 72: 340-343.
2. Dissing, J., et al. 1987. Human red cell acid phosphatase (ACP1): evidence for differences in the primary structure of the two isozymes encoded by the ACP1*B allele. *Biochem. Genet.* 25: 919-927.
3. Dissing, J., et al. 1991. Human red cell acid phosphatase (ACP1). The amino acid sequence of the two isozymes Bf and Bs encoded by the ACP1*B allele. *J. Biol. Chem.* 266: 20619-20625.
4. Wo, Y.Y., et al. 1992. Sequencing, cloning, and expression of human red cell-type acid phosphatase, a cytoplasmic phosphotyrosyl protein phosphatase. *J. Biol. Chem.* 267: 10856-10865.

CHROMOSOMAL LOCATION

Genetic locus: Acp1 (mouse) mapping to 12 A2.

PRODUCT

ACP1 shRNA (m) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200 μ l frozen stock containing 1.0×10^6 infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see ACP1 siRNA (m): sc-44359 and ACP1 shRNA Plasmid (m): sc-44359-SH as alternate gene silencing products.

STORAGE

Store lentiviral particles at -80°C . Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4°C for up to one week. Avoid repeated freeze thaw cycles.

PROTOCOLS

See our web site at www.scbt.com or our catalog for detailed protocols and support products.

APPLICATIONS

ACP1 shRNA (m) Lentiviral Particles is recommended for the inhibition of ACP1 expression in mouse cells.

SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200 μ l frozen viral stock containing 1.0×10^6 infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

GENE EXPRESSION MONITORING

ACP1 (D-3): sc-390190 is recommended as a control antibody for monitoring of ACP1 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support (secondary) reagents are recommended: 1) Western Blotting: use goat anti-mouse IgG-HRP: sc-2005 (dilution range: 1:2000-1:32,000) or Cruz Marker™ compatible goat anti-mouse IgG-HRP: sc-2031 (dilution range: 1:2000-1:5000), Cruz Marker™ Molecular Weight Standards: sc-2035, TBS Blotto A Blocking Reagent: sc-2333 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use goat anti-mouse IgG-FITC: sc-2010 (dilution range: 1:100-1:400) or goat anti-mouse IgG-TR: sc-2781 (dilution range: 1:100-1:400) with UltraCruz™ Mounting Medium: sc-24941.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ACP1 gene expression knockdown using RT-PCR Primer: ACP1 (m)-PR: sc-44359-PR (20 μ l). Annealing temperature for the primers should be $55-60^\circ\text{C}$ and the extension temperature should be $68-72^\circ\text{C}$.

BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.