



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

CaSR siRNA (h): sc-44373

BACKGROUND

Extracellular calcium-sensing receptor (CaSR), also designated parathyroid cell calcium-sensing receptor, is an integral membrane protein that belongs to the G protein-coupled receptor 3 family. CaSR is involved in maintaining a stable calcium concentration by acting as a sensor of the extracellular calcium levels for the parathyroid and kidney. Its activity is mediated by a G protein which activates a phosphatidylinositol-calcium second messenger system. Defects that activate CaSR cause autosomal dominant hypocalcemia, whereas mutations that inactivate the protein cause familial hypocalciuric hypercalcemia. CaSR is expressed mainly in kidney, and is also expressed in intestine, placenta and brain.

CHROMOSOMAL LOCATION

Genetic locus: CASR (human) mapping to 3q21.1.

PRODUCT

CaSR siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CaSR shRNA Plasmid (h): sc-44373-SH and CaSR shRNA (h) Lentiviral Particles: sc-44373-V as alternate gene silencing products.

For independent verification of CaSR (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-44373A, sc-44373B and sc-44373C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

CaSR siRNA (h) is recommended for the inhibition of CaSR expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

CaSR (6D4): sc-47741 is recommended as a control antibody for monitoring of CaSR gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CaSR gene expression knockdown using RT-PCR Primer: CaSR (h)-PR: sc-44373-PR (20 μ l, 443 bp). Annealing temperature for the primers should be $55-60^{\circ}$ C and the extension temperature should be $68-72^{\circ}$ C.

SELECT PRODUCT CITATIONS

1. Molostvov, G., et al. 2008. Extracellular calcium-sensing receptor mediated signalling is involved in human vascular smooth muscle cell proliferation and apoptosis. *Cell. Physiol. Biochem.* 22: 413-422.
2. Li, H., et al. 2009. The calcimimetic R-568 induces apoptotic cell death in prostate cancer cells. *J. Exp. Clin. Cancer Res.* 28: 100.
3. Molostvov, G., et al. 2015. Arterial expression of the calcium-sensing receptor is maintained by physiological pulsation and protects against calcification. *PLoS ONE* 10: e0138833.
4. D'Espessailles, A., et al. 2018. Calcium-sensing receptor activates the NLRP3 inflammasome in LS14 preadipocytes mediated by ERK1/2 signaling. *J. Cell. Physiol.* 233: 6232-6240.
5. Shin, S., et al. 2019. L-ornithine activates Ca²⁺ signaling to exert its protective function on human proximal tubular cells. *Cell. Signal.* 67: 109484.
6. Liu, L., et al. 2020. CaSR induces osteoclast differentiation and promotes bone metastasis in lung adenocarcinoma. *Front. Oncol.* 10: 305.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.